







Exercise therapy for chronic symptomatic peripheral artery disease

A clinical consensus document of the European Society of Cardiology Working Group on Aorta and Peripheral Vascular Diseases in collaboration with the European Society of Vascular Medicine and the European Society for Vascular Surgery

Lucia Mazzolai ^{1*}†, Jill Belch^{2†}, Maarit Venermo ^{3,4†}, Victor Aboyans ^{5,6}, Marianne Brodmann⁷, Alessandra Bura-Rivière⁸, Sebastien Debus⁹, Christine Espinola-Klein ¹⁰, Amy E. Harwood¹¹, John A. Hawley¹², Stefano Lanzi ¹, Juraj Madarič^{13,14}, Guillaume Mahé^{15,16}, Davide Malatesta¹⁷, Oliver Schlager ¹⁸, Arno Schmidt-Trucksäss¹⁹, Chris Seenan²⁰, Henrik Sillesen^{21,22}, Garry A. Tew²³, and Adriana Visonà²⁴

¹Angiology Department, Lausanne University Hospital, University of Lausanne, Ch. de Mont-Paisible 18, Lausanne 1011, Switzerland; ²Institute of Cardiovascular Research, University of Dundee, Ninewells Hospital and Medical School, Dundee, UK; ³Department of Vascular Surgery, Abdominal Center, Helsinki University Hospital, Helsinki, Finland; ⁴University of Helsinki, Helsinki, Finland; ⁵Department of Cardiology, Dupuytren-2 University Hospital, France; ⁶EpiMaCT, INSERM 1094/IRD270, Limoges University, Limoges, France; ⁷Division of Angiology, Department of Internal Medicine, Medical University, Graz, Austria; ⁸Department of Vascular Medicine, Toulouse University Hospital, Toulouse, France; ⁹Department of Vascular Medicine, Vascular Surgery–Angiology–Endovascular Therapy, University of Hamburg-Eppendorf, Hamburg, Germany; ¹⁰Center of Cardiology, Department of Cardiology III-Angiology, University Medical Center of the Johannes Gutenberg-University Mainz, Mainz, Germany; ¹¹Department for Sport and Exercise Sciences, Manchester Metropolitan University, Manchester, UK; ¹²Exercise and Nutrition Research Programme, Mary MacKillop Institute for Health Research, Australian Catholic University, Melbourne, VIC, Australia; ¹³Department of Angiology, Comenius University, Bratislava, Slovakia; ¹⁴National Institute of Cardiovascular Diseases, Bratislava, Slovakia; ¹⁵Vascular Medicine Unit, Centre Hospitalier Universitaire de Rennes, Rennes, France; ¹⁶INSERM CIC 1414, Université de Rennes, Rennes, France; ¹⁷Institute of Sport Sciences, University of Lausanne, Lausanne, Switzerland; ¹⁸Division of Angiology, Department of Medicine II, Medical University of Vienna, Vienna, Austria; ¹⁹Division of Sport and Exercise Medicine, Department of Sport, Exercise and Health, University of Basel, Basel, Switzerland; ²⁰School of Health and Life Sciences, Glasgow Caledonian University, Glasgow, UK; ²¹Department of Vascular Surgery, Rigshospitalet, Copenhagen, Denmark; ²²Department of Clinical Medicine, University of Copenhagen, Denmark; ²³Institute for Health and Care Improvement, York St John University, York, UK; and ²⁴Angiology Unit, Ospedale Castelfranco Veneto, Castelfranco Veneto, Italy

Abstract


All guidelines worldwide strongly recommend exercise as a pillar of the management of patients affected by lower extremity peripheral artery disease (PAD). Exercise therapy in this setting presents different modalities, and a structured programme provides optimal results. This clinical consensus paper is intended for clinicians to promote and assist for the set-up of comprehensive exercise programmes to best advice in patients with symptomatic chronic PAD. Different exercise training protocols specific for patients with PAD are presented. Data on patient assessment and outcome measures are narratively described based on the current best evidence. The document ends by highlighting disparities in access to supervised exercise programmes across Europe and the series of gaps for evidence requiring further research.

* Corresponding author. Tel: +41 021 314 07 68, Email: lucia.mazzolai@chuv.ch

† Shared first authors.

This article has been co-published with permission in European Heart Journal, Vasa – European Journal of Vascular Medicine and European Journal of Vascular and Endovascular Surgery. © 2024 the European Society of Cardiology, the European Society of Vascular Medicine, and the European Society for Vascular Surgery. Published by Oxford University Press on behalf of the European Society of Cardiology, by Hogrefe AG on behalf of European Society of Vascular Medicine and by Elsevier B.V on behalf of European Society for Vascular Surgery. All rights reserved. The articles are identical except for minor stylistic and spelling differences in keeping with each journal's style. Either citation can be used when citing this article.

Graphical Abstract



Graphical summary of the exercise training approaches in patients with peripheral artery disease.

Keywords

Vascular rehabilitation • Quality of life • Intermittent claudication • Exercise training • Physical activity

Introduction

Physical activity, including regular exercise, is one of the pillars of cardiovascular (CV) health and a major component of management of patients with most CV diseases (CVD). In 2020, the European Society of Cardiology (ESC) issued a guideline document addressing the main aspects of exercise therapy and sports practice for cardiac diseases.¹

In this consensus document, the acronym PAD will be used to indicate lower extremity peripheral artery disease. Peripheral artery disease is one of the most prevalent clinical presentations of atherosclerotic disease, affecting ~237 million people worldwide.² The first symptoms of PAD are usually related to walking impairment, and the 2017 ESC/European Society for Vascular Surgery (ESVS) guidelines on the management of PAD underscore the importance of exercise therapy, preferably supervised, for the management of patients with intermittent claudication (IC).³ Similarly, the 2019 PAD guidelines of the European Society of Vascular Medicine (ESVM) encourage structured exercise for symptomatic PAD patients.⁴ However, none of the aforementioned documents provided in-depth guidance for exercise therapy in this specific setting.

To address this gap, the ESC Working Group on Aorta and Peripheral Vascular Diseases, the ESVM, and the ESVS joined in a

collaborative effort aiming to provide a roadmap and guidance for the set-up and implementation of exercise therapy programmes for patients with PAD (Graphical Abstract).

Consensus statements

- (1) In patients with PAD and exercise-induced limb symptoms due to vascular origin, supervised exercise programmes should be the first-line treatment modalities.
- (2) In patients with PAD undergoing revascularization, supervised exercise programmes should be included as adjuvant therapy.
- (3) Supervised exercise programmes should ideally be coordinated by vascular physicians, and sessions should be ideally supervised by clinical exercise physiologists or physiotherapists.
- (4) Prior to exercise training initiation, complete medical history and examination and screening for contraindications should be investigated.
- (5) Measures of walking ability, functional status, and quality of life should be assessed at the beginning and end of the programme to determine the patient's response to exercise training. Clinical outcomes and patient experience should also be documented.




Figure 1 Pathophysiology of limb symptoms in peripheral artery disease

- (6) Walking training (overground, pole striding, treadmill) should be proposed as first-line exercise modality. When walking is not an option, alternative training modalities (resistance and strength training, arm-cranking, cycling, combinations of exercise) should be performed.
- (7) The training frequency should be at least three times per week.
- (8) The training session duration should last a minimum of 30 min.
- (9) The training programme duration should last a minimum of 3 months.
- (10) Both claudication pain (A) and exercise intensity [B, based on common training intensity measures such as heart rate (HR) or the rate of perceived exertion (RPE) on Borg's scale] should be evaluated during training sessions:
 - (A) The current consensus is that patients should exercise to moderate–high claudication pain based on strong evidence. However, some trials have recently demonstrated improvement in walking ability using a low or no pain approach. As claudication pain is a commonly cited barrier to exercise, universally prescribing high-pain exercise may lead to poor uptake of, and adherence to, exercise training programmes. A more flexible approach to exercise prescription may therefore be required, considering the patient's needs and preferences and what might achieve a high level of (long-term) adherence.
 - (B) Following a 'lead-in period' of low-to-moderate exercise intensity, a gradual progression to vigorous/high exercise intensity may be proposed if well tolerated by the patient.
- (11) If supervised exercise is not available or feasible, a structured community- or home-based exercise programme that includes behaviour change techniques should be proposed.
- (12) Supervised exercise programmes should include structured education and counselling on CVD and PAD risk factor reduction. Smoking cessation should be a cornerstone of risk factor counselling.
- (13) Following initial exercise training (supervised or home-based), patients are encouraged to sustain lifelong and high levels of regular physical activity.

Pathophysiology of intermittent claudication and functional impairment

Intermittent claudication is characterized by exertional leg pain limiting walking ability.^{5–7} Peripheral artery disease induces a wide range of exercise-related symptoms experienced by nearly half of the PAD population.⁸ The classical IC symptomatology was first defined as calf pain, discomfort, or fatigue appearing during exercise and forcing the patient to stop.⁹ Typically, IC is relieved within 2–5 min after discontinuation of exertion.⁹ Apart from this typical symptom, it is now admitted that some patients with PAD may present atypical exercise-induced limb symptoms.¹⁰ These may be localized in lower limb muscles other than calves; may be present at rest; may be described by patients as 'burning', 'compressive' feeling, or just 'fatigue' without pain; and may mimic limb pain due to spinal stenosis. Exercise-induced limb symptoms in PAD are caused by a metabolic mismatch between oxygen demand and supply.⁵ The mismatch is linked to the reduction of the arterial lumen by the atherosclerosis process, but it also induces cellular and metabolic disorders that contribute to the functional impairment.¹¹ Mechanisms of exercise-induced symptoms are multifactorial among which nociceptive pain,¹² nerve dysfunction,¹³ and skeletal muscle abnormalities¹¹ are suggested.

Potential mechanistic drivers of exertional limb symptoms in addition to arterial obstruction and reduced perfusion include inflammation, vascular dysfunction, reduced microvascular flow, impaired angiogenesis, and altered skeletal muscle function^{14–16} (Figure 1). A healthy vascular endothelium produces several vasodilator substances, including nitric oxide (NO), which has pluripotent vascular benefits such as platelet inhibition, smooth muscle cell proliferation inhibition, leukocyte adhesion prevention, and angiogenesis induction. Diminished NO bioactivity in the lower limbs prevents increased blood flow with exercise.¹¹ Vascular dysfunction may also exacerbate the vasoconstrictive effects of catecholamines and limit flow-mediated dilation.^{17–20} Inadequate angiogenesis and collateral vessel formation may potentiate limb ischaemia and serve as a mechanism driving functional impairment.²¹ Skeletal muscle ischaemia may drive local inflammation, exacerbating symptoms and altering muscle metabolism.^{22–24}

Patients with PAD present impaired walking endurance,²⁵ slower walking velocity,^{26–28} gait abnormalities,^{26,27,29–31} poorer muscle strength,³²

and poorer balance^{33,34} compared with individuals without PAD. They may also reduce their walking activity and total activity to avoid leg symptoms,³⁵ and studies have shown a functional decline occurring over time.^{25,28,36}

Vascular and functional assessment in peripheral artery disease

Vascular assessment

General assessment of CV risk factors should be performed prior to exercise training rehabilitation to improve preventive measures and reach preventive goals. Ankle-brachial index (ABI) should be assessed before starting a training programme to detect and diagnose PAD and assess disease severity (Figure 2).³ The measurement of ABI after exercise is also important to further detect ankle pressure drop, as some patients may have leg symptoms on exercise, while ABI can be ≥ 0.91 at rest. A post-exercise ankle systolic blood pressure drop >30 mmHg or a post-exercise ABI decrease $>20\%$ should be considered for PAD diagnosis.³⁷ In patients with media calcinosis (e.g. in patients with diabetes or chronic kidney disease) measurement of ABI might not be possible because the arteries cannot be compressed by the cuff. In these cases, toe-brachial index (TBI) can be used as alternative assessment (the pathological threshold usually retained is <0.70).³

Walking distance assessment

Walking distance is considered an important clinical outcome both for patients and clinicians. Standardized exercise testing should be used for assessment of functional impairment in patients with PAD (Figure 2).

Treadmill assessment

Treadmill testing should be performed with patients familiarized to the treadmill and under reproducible conditions (i.e. avoiding exercise and alcohol prior to assessment). Patients should be asked to walk until maximal levels of pain, lightly holding or not holding onto the treadmill. If the tests are stopped for reasons other than leg pain, then this should be recorded. Patients are asked to indicate the claudication pain score they reached during walking, especially the point at which pain begins, and recovery based on a five-point scale (0, no pain; 1, onset of pain; 2, mild pain; 3, moderate pain; 4, severe/maximal pain).³⁸ Common treadmill protocols include constant-load (single-stage) or graded exercise testing.^{39,40} The latter is performed at constant speed varying the slope of the treadmill. Established graded protocols include the Gardner/Skinner (3.2 km/h and a 2% increase in slope every 2 min) or the Hiatt protocol (3.2 km/h and an increase in slope of 3.5% every 3 min). Constant-load treadmill tests are performed at a fixed speed of 2–4 km/h and fixed gradient of 10%–12%. Constant-load protocols have poorer reliability both for pain-free walking distance (PFWD) and maximal walking distance (MWD) compared with graded protocols (coefficient of variance 30% and 45%, respectively).^{41,42} Treadmill tests have limitations including learning effect during repeated evaluations. Also, some patients are unable or are unwilling to perform a treadmill test, mainly due to balance impairment or limited walking abilities.

Six-minute walk test

The six-minute walk test (6MWT) is performed along a flat corridor with a length of 30 m with turning points marked by a cone. Patients are asked to walk self-paced for the full duration and may stop and rest at any point in the test.⁴³ The total distance walked is measured and reported as the six-minute walking distance (6MWD).⁴³ Any encouragement given/

phrases used should be the same for every test performed to ensure test-retest reliability.⁴³ Further, there may be a learning effect so it is recommended that the best out of two walks is recorded or the first test discounted.⁴⁴ Although treadmill-based exercise tests can establish maximum walking capacity, there may be a poor correlation between treadmill outcomes, habitual walking, and self-reported walking distance.⁴⁵ On the other hand, compared with treadmill test, the 6MWT has been shown to better represent daily life walking in patients with PAD.⁴⁶ The 6MWT is a well-validated and low-cost test. It has good reliability, with a correlation coefficient of 0.90 ($P < .001$) and a coefficient of variation of 8.9% with testing performed 1 to 2 weeks apart.⁴⁷ Changes in the 6MWT can be used to predict mortality and mobility loss in patients with PAD.^{7,48} The minimal detectable changes (i.e. the statistical detectability of change beyond measurement error) in the 6MWT are represented by a change >46 m.⁴⁹ The minimal clinically important difference (i.e. the clinical relevance or importance of the observed change from the patient's perspective) in the 6MWT in patients with PAD is represented by an improvement of 8⁵⁰ or 9 m⁵¹ for small changes and 20⁵⁰ or 38 m⁵¹ for large changes.

Connected devices

A measure of 'real-life' walking performances may be performed by the use of global positioning systems (GPS) or commercially available devices such as activity trackers, smart watches, and phones.⁵² Research has shown that GPS recorders have good accuracy and reliability when compared with known distances walked,^{53,54} and measurement of step counts with mobile phones has been shown to be highly reliable even at low walking speeds.⁵⁵ Further, GPS recorded walking distances correlate well with treadmill walking distances.⁵⁶ Patients should be able to note the initial onset of claudication pain and the MWD either in total or between bouts of walking using the GPS system.

Muscle strength assessment

The presence of PAD is associated with impaired lower extremity muscle strength and function,⁵⁷ which is associated with high prevalence of frailty and sarcopenia.⁵⁸ Muscle strength and function should therefore be assessed before and after supervised exercise training (SET; Figure 2). There is heterogeneity in how muscle strength and function are assessed. Muscle isokinetic strength and endurance can be assessed via isokinetic dynamometry, which is a chair device that patients sit on and the specific joint is tested in an appropriate position with the dynamometer attached to the limb. Patients push against the dynamometer as it provides resistance to maintain a set speed. Isokinetic dynamometry has demonstrated good reliability at the ankle (reliability coefficients ranging from 0.77 to 0.96).⁵⁹ Testing can be done in various joints, including the ankle, knee, and hip, in various planes such as extension and flexion. As isokinetic dynamometry assessment includes specialized equipment, it may not be practical or convenient to assess patients using this device. As an alternative, the short physical performance battery (SPPB) which includes a 4 m walk test, a sit-to-stand chair test, and a standing balance test should be used.⁶⁰ A recent study showed that the sit-to-stand is a validated test to estimate muscle power in patients with symptomatic PAD.⁶¹ Interestingly, muscle power assessed by the sit-to-stand test was related to overall functional performance prior and following SET.⁶¹

Self-reported functional impairment and quality of life assessment

In addition to objective assessment of functional impairment, a subjective (self-reported) evaluation of walking abilities and health-related




Figure 2 Algorithm of chronic symptomatic patients with PAD with indication for exercise treatment. PAD, peripheral artery disease; SPPB, short physical performance battery; BMT, best medical treatment [including pharmacological and non-pharmacological (lifestyle changes and exercise) approach]; DUS, duplex ultrasound; SF-36, short-form health 36 questionnaire; WIQ, Walking Impairment Questionnaire; Vascu-QoL6, Vascular Quality of Life Questionnaire-6

Table 1 Self-reported evaluation of walking ability and health-related quality of life in patients with peripheral artery disease

Questionnaire name	Type (functional/QOL)	Domains tested
EQ-5D	General	Mobility, self-care, usual activity, pain/discomfort, and anxiety/depression
WHOQOL	General	Physical health, psychological health, social relationships, and environment
SEIQoL	General	Five dimensions chosen by the patient
VascuQOL	PAD-specific	Pain, symptoms, activities, and social and emotional well-being
ICQ	PAD-specific	Walking distance, walking speed, and stair climbing
PADQOL	PAD-specific	Social relationship and interaction, self-concept and feelings, symptoms and limitations in physical functioning, fear and uncertainty, and positive adaptation while living with PAD
SF-36	General	Physical function, bodily pain, general health, mental health, vitality, emotional well-being, and social functioning
NHP	General	Energy, emotional reaction, sleep, pain, social isolation, and physical mobility
Peripheral Artery Questionnaire (PAQ)	PAD-specific	Physical limitations, symptoms, social function, treatment satisfaction, and quality of life
Walking Impaired Questionnaire (WIQ)	PAD-specific	Physical limitations and symptoms
Walking Estimated-Limitation Calculated by History (WELCH)	PAD-specific	Physical limitations and symptoms

Table 2 Absolute contraindications to exercise training

Acute coronary syndrome (within 2 days)
Unstable cardiac disease on interview/examination
Uncompensated heart failure
Acute thrombophlebitis or recent embolism (pulmonary or systemic)
Active endocarditis
Acute myocarditis or pericarditis
Acute aortic dissection
Symptomatic severe aortic stenosis
Acute systemic illness or fever
Uncontrolled hypertension (≥ 180 mmHg systolic or ≥ 110 mmHg diastolic blood pressure at rest)
Uncontrolled sinus tachycardiac (resting heart rate > 120 beats/min)
Third-degree atrioventricular block without pacemaker
Uncontrolled diabetes mellitus
Orthostatic drop in blood pressure (> 20 mmHg) with symptoms

Adapted from the American College of Sports Medicine (2022) ACSM's guidelines for exercise testing and prescription. 11th ed Guidelines for exercise testing and prescription. Philadelphia, PA: Lippincott Williams & Wilkins.

quality of life (HRQoL) should be incorporated to have a complete assessment of the functional status of the patient (Figure 2).^{62–64} Following exercise interventions, assessing HRQoL is usually used to determine if an

objective improvement in functional performance is also perceived by the patients in their daily life. Table 1 reports the most used subjective tools used for walking ability and HRQoL assessment in patients with PAD. Trials used a wide variety of questionnaires of patient-reported outcome measurements (PROMs).^{62–64} The most used are the short-form health 36 (SF-36), a generic questionnaire including physical and mental items related to health, and the Walking Impairment Questionnaire (WIQ), a PAD-specific questionnaire focusing on PAD and functional limitations. Studies have shown that HRQoL burden is greater in magnitude in patients with both PAD and CVD than with CVD alone.⁶⁵ In the PARTNERS study, the SF-36 Physical Component Summary of the combined PAD-other-CVD group was 46.3 ± 1.2 compared with 55.5 ± 1.1 in the other-CVD group.⁶⁵ Cross-sectional studies show that in patients with PAD the degree of difficulty in walking distance and stair climbing are significantly related to HRQoL.⁶⁶ The ESVS VASCUNET and the International Consortium of Vascular Registries consensus statement recommended the Vascular Quality of Life Questionnaire-6 (Vascu-QoL6) as a primary assessment of PROMs in patients with symptomatic PAD.⁶²

Greater amounts of physical activity are associated with higher ratings of both perceived health and HRQoL, correlating with objective health outcomes and life expectancy.⁶⁷ One of the most important factors linked to both subjective and objective health, across both cognitive and physical domains, is physical activity.⁶⁸

Exercise therapy in patients with peripheral artery disease

Screening prior to exercise training participation

All patients should be medically screened before SET programme initiation (Figure 2). It is suggested to include a complete medical history and examination.³⁸ Patients with contraindications to exercise training (Table 2) should be excluded from SET until the relevant condition

Table 3 Conditions needing a cardiological screening before exercise training participation

History of documented coronary artery disease
History of documented major arrhythmia and atrial fibrillation
History of documented congenital heart disease
Any clinical sign or electrocardiogram suspicion for cardiac disease

stabilizes or is successfully treated. For patients with current or prior symptomatic cardiac disease (Table 3), we recommend that they are referred for cardiology work-up, including an exercise test to assess for evidence of exercise-induced coronary ischaemia, to identify if additional treatment for cardiac disease is required before proceeding with SET. Comorbidities (such as neurological and orthopaedic diseases leading to gait abnormalities) should be documented and considered for how they may limit SET programme participation feasibility. After SET programme initiation, patients should continue to be closely monitored for changes in health status (e.g. any symptom or situation which may suspect undiagnosed/incident cardiac condition, ischaemic limb pain at rest, and toe or foot wounds) that might necessitate interruption of the programme, at least temporarily.

Supervised exercise training

Supervised exercise training is considered among first-line therapies for patients with chronic and symptomatic PAD (Figure 2).^{3,64,69,70} Supervised exercise training is safe and is usually conducted in the hospital setting.⁷¹ Over the past 60 years, many trials have reported the effectiveness of SET on walking distances in these patients.^{72,73} The most recent Cochrane meta-analysis showed that SET improves PFWD (82 m; 95% CI: 72–92) and MWD (120 m; 95% CI 51–190).⁷⁴ Similar findings were observed in another meta-analysis [PFWD: 128 m (95% CI: 92–165); MWD: 180 m (95% CI: 130–230)].⁷⁵ Although less well investigated or usually reported as a secondary outcome, SET also improved functional status, gait pattern, self-reported walking ability, and quality of life.^{64,74,76–82} It is interesting to note that cardiac rehabilitation programmes also increase walking distance, HRQoL, and physical activity in patients with symptomatic PAD, suggesting that other types of rehabilitation than SET may also be useful.⁸³ Finally, some vasoactive drugs such as cilostazol (phosphodiesterase type 3 inhibitor), pentoxifylline (xanthine derivative), bosentan, sildenafil, and others are claimed to increase walking capacity in patients with PAD.^{84–87} However, the objective documentation of their effect is very limited to draw extensive conclusions.^{84,88} More studies are needed to confirm additive effect of drug therapies to supervised exercise.

Training modalities

There are different types of exercise training for patients with PAD, but the common aim is to improve walking capacity and reduce symptoms. In addition, exercise should aim to improve balance and muscle strength to promote independence and a reduced risk of falling in the long term.³³ Treadmill and overground walking are the most common and recommended training modalities in patients with IC (Figure 2).^{64,70} However, due to severe exercise-induced ischaemia, low pain tolerance, the risk of falling, and/or other comorbidities, some patients are unwilling or unable to perform walking sessions. In addition to walking

training, there are several other forms of training that are used, although much less frequently, in the rehabilitation of patients with PAD. A recent meta-analysis reported that other non-walking training modes are also effective as traditional walking training in improving walking performance, whereas there was no clear evidence for changes in quality of life following exercise interventions. However, the authors concluded that the certainty of this evidence was judged to be low.⁸⁹ Different training modes include strength training of large muscle groups,^{90,91} cycling,⁹² pole striding,^{93,94} multimodal training,^{76,77,95–98} and training with an arm-crank ergometer.^{99,100} The beneficial effect of these training modalities can usually be described as large and even reach those of typical walking training.¹⁰¹ However, the PFWD and the MWD have the tendency to be higher with walking training than with strength training when all studies are considered.⁸⁹ In contrast, self-reported ability to climb stairs (assessed by the WIQ) is more improved following strength training (29.2% vs. 43.8% after 6 months) compared with walking training on the treadmill (39.6% vs. 43.8% after 6 months).¹⁰² Therefore, when walking is not an option, alternative training modalities might also be effective. These training modalities also elicit lower or no pain during exertion compared with walking, which might lead to higher rates of adherence.

Training frequency

Based on a previous meta-analysis, and shared by most of the studies and guidelines, the training frequency associated with greater improvements in walking distance is at least 3 times per week.^{103,104}

Training duration

Identifying an optimal training duration is difficult to elucidate, mainly due to differences in training modalities, frequencies, and intensities among studies. Current guidelines reported that optimal training duration ranges between 12 and 24 weeks.^{64,70,103} The optimal training session duration has not been widely investigated. Additionally, in most studies, the total session duration is usually reported without specifying the actual time spent exercising. The literature shows that exercise sessions lasting 30–60 min were the most effective to improve walking performance.^{103,104}

Training intensity

In most studies, no clear distinction is made between symptom intensity (claudication pain scale) and exercise training intensity (based on HR, oxygen uptake (VO₂), or RPE on Borg's scale: 6, 'very very light'; 20, 'maximal effort') to monitor the exercise therapy. The Borg scale is a subjective assessment tool used to measure an individual's perceived exertion or effort during physical activity. The scale assigns a numerical rating ranging from 6 to 20 to indicate the intensity of exertion experienced by the person.¹⁰⁵

First, the majority of trials used claudication pain severity to provide guidance during the training sessions. In PAD research, the claudication pain scale, an ordinal scale from 0 (no pain) to 4 (severe/maximal pain), is the most commonly used tool. A distinction is made between walking training with and without muscle pain caused by ischaemia. With regard to claudication pain intensity, international guidelines are heterogeneous.^{38,64,70} The UK NICE guideline encourages patients to exercise to the point of maximal pain,¹⁰⁶ the American Heart Association guideline recommends moderate-to-moderate/severe claudication pain as tolerated,⁶⁴ while the Australian guideline does not specify pain intensity for exercise dosage.¹⁰⁷ Based on strong evidence,^{64,73–75,104} the current consensus is that patients should exercise to moderate–high

claudication pain to improve walking performance. Also, 1 year home-based walking training performed at high-intensity pain has been found to be more effective than walking training performed at low intensity for improving walking and functional performance in patients with PAD.^{108,109} These findings indicate that claudication pain intensity may be a key factor for walking improvement in these individuals. In contrast, others have reported that improvements in walking performance may be obtained with less severe claudication pain during exertion.¹⁰¹ According with recent findings, walking training with pain is not clearly superior to walking training without pain regarding changes in walking distances.^{110–113} It may be assumed that walking training with moderate, low, or no pain is associated with higher compliance and possibly long-term maintenance of training or change in activity behaviour.¹¹³ This indicates that a more flexible approach to exercise prescription may therefore be required, considering the patient's needs and preferences, and what might achieve a high level of (long-term) adherence. Larger studies with a higher number of cases and longer duration, taking compliance into account, are needed for a conclusive statement.¹¹⁴

Second, the optimal no/low pain-based exercise training intensity is understudied in this population. Indeed, it is interesting to note that the claudication pain severity does not necessarily rely on common measures of exercise intensity.^{78,115} For example, when performing vigorous-intensity exercise, some patients may experience moderate-to-severe claudication pain, whereas others, low levels of claudication only. Assuming that exercise intensity is a cornerstone determinant of physiological response to training,¹¹⁶ monitoring claudication pain only is limiting and prevents accurate comparison of exercise effectiveness in patients with PAD. This may also explain the large variability in the magnitude of improvements following exercise interventions.^{64,103} Fassora et al.⁷⁸ recently reported that both training modality and exercise intensity should be considered when looking for the best results in walking performance and cardiorespiratory fitness. Notably, these results showed that walking at vigorous intensity (%HR_{peak}: 77–95, %VO_{2peak}: 64–90, RPE: ≥ 14¹¹⁶) induced the greatest improvement in MWD, while cycling and other non-walking modalities performed at vigorous intensity elicited the greatest improvements in cardiorespiratory fitness.⁷⁸ These findings suggest that both walking and cardiorespiratory capacities are desirable outcomes but that they need different exercise therapy programmes.⁷⁸ It is however important to note that training programmes should start with a lead-in period performed at low-to-moderate exercise intensity and, if tolerated, gradually progressed to vigorous exercise intensity. This approach may allow to determine the patient's exercise response and tolerance, reducing the risk of complications.

The monitoring of the exercise intensity during a resistance training programme is mediated by the percentage of the one repetition maximum (1RM).¹¹⁷ The determination of the 1RM plays a key role to objectively set an individualized resistance-based programme.¹¹⁷ Compared with a direct assessment of the 1RM, the multiple RM assessment (such as 10RM, the maximum weight a person can lift for 10 repetitions) is considered to be a safe and well-tolerated approach to evaluate muscle strength for a given muscle group in patients with CVDs.¹¹⁷ Following the multiple RM test, different prediction equations are available to estimate the 1RM.¹¹⁸ As also used in the cardiac rehabilitation, a target exercise intensity of 30%–70% of 1RM for the upper body and 40%–80% of 1RM for the lower body should be considered.¹¹⁸ Exercise intensity should be progressively increased to determine the patient's exercise response and exercise tolerance. It has been shown that resistance training improves walking performance

and muscular strength in patients with PAD.¹¹⁹ Notably, high intensity (i.e. 80% 1RM) induces the best improvements in walking performance when compared with low-to-moderate (i.e. <50% 1RM) strength training intensity in these patients.^{90,119}

Table 4 summarizes the main exercise prescription recommendations with some practical applications.

Home-based exercise training

In comparison with patients not undergoing exercise training, a home-based training (HBT) strategy resulted in a non-significant increase of MWD in a recent meta-analysis (mean difference: 136 m; 95% CI: –2 to 273 m; $P = .05$).¹²⁰ When comparing HBT with basic exercise advice, no improvement of MWD was observed in patients following a HBT strategy (mean difference: 39 m; 95% CI: –123.1 to 201.1 m; $P = .64$).¹²⁰ Regarding PFWD, HBT led to a greater increase than exercise advice did (mean difference: 64.5 m; 95% CI: 14.1–114.8 m; $P = .01$).¹²⁰ In comparison with HBT, SET was more effective in improving MWD (mean difference: 139 m; 95% CI: 45–232 m; $P = .004$) and PFWD (mean difference: 84 m; 95% CI: 25–143 m; $P = .005$).¹²⁰

Considering the effect of monitoring in HBT, no difference in the change of MWD and PFWD was observed between monitored HBT and SET (mean difference in MWD: 8 m; 95% CI: –81 to 97 m; $P = .86$; mean difference in PFWD: 43 m; 95% CI: –29 to 114 m; $P = .24$).¹²⁰ The equality in training efficacy of monitored HBT and SET emphasizes the role of monitoring in HBT programmes. Apart from regular on-site visits or phone calls, activity diaries or log books have been used for HBT monitoring.¹²⁰ Additional tools for self-monitoring, such as wrist-worn activity trackers with smartwatch-like functions or smartphone accelerometer applications, have been assessed; however, it still needs to be clarified, which modality is most appropriate.⁵⁵

The effect of training on patients' daily physical activity was assessed by several studies implementing pedometer and accelerometer measurements. A network meta-analysis demonstrated improvements of daily physical activity in HBT to a similar extent as it was observed in patients undergoing SET.¹²¹

Focusing on quality of life, most studies reported improvements in patients undergoing HBT.¹²⁰ In comparison with SET, improvements of individual SF-36 measures (pain and social functioning) and WIQ measures (distance) were less pronounced in patients undergoing HBT.¹²⁰ In addition, HBT improves measures of self-efficacy for walking, satisfaction with functioning, pain acceptance, and social functioning in patients with claudication.¹²² Follow-up data of patients who had undergone HBT suggest sustained improvements in measures of quality of life and functional and walking capacity after termination of the active training intervention.^{123,124}

Safety of HBT was analysed in a systematic review including 27 studies, which reported a cardiac event rate of 1 per 49 270 and a non-cardiac event rate of 1 per 147 810.¹²⁵ Event rates of HBT were lower than event rates reported for SET (HBT vs. SET: cardiac 1:49 270 vs. 1:13 788; non-cardiac 1:147 810 vs. 1:41 363).¹²⁵ Regarding overall mortality, retrospective data suggest a reduction of long-term mortality in patients undergoing HBT.¹²⁶ Comparing HBT with SET, overall mortality rates do not differ between patients undergoing HBT and patients following a SET programme.¹²⁷ The results of the reported meta-analyses and reviews should be viewed with caution according to a moderate to low quality of evidence.^{120,127,128} Due to the limited availability and utilization of SET programmes, HBT programmes can be used as a valid alternative training modality for patients with IC.^{129–132}

Table 4 Training specificity and practical applications

Training modality	Training frequency	Training duration	Claudication pain intensity	Exercise intensity ^a	Example protocols
Walking (treadmill or overground)	At least 3x per week	<p><i>Session's duration</i></p> <ul style="list-style-type: none"> Start at 10–15 min of actual exercise time. Increase progressively up to 30–60 min of actual exercise time (including warm-up). <p><i>Programme duration</i></p> <ul style="list-style-type: none"> At least 12 weeks. Following initial exercise training, patients are encouraged to sustain lifelong and high levels of regular physical activity. 	<ul style="list-style-type: none"> Moderate–high Mild Pain-free 	<p><i>Low-to-moderate</i></p> <ul style="list-style-type: none"> HR_{peak}: ≤ 76% RPE: ≤ 13 <p><i>Vigorous</i></p> <ul style="list-style-type: none"> HR_{peak}: 77%–95% RPE: ≥ 14 	<p><i>For people who are able and willing to walk at moderate–high pain intensity</i></p> <ul style="list-style-type: none"> Walk at a speed and/or grade that induces moderate–high (3–4 on the claudication pain scale) claudication pain intensity Rest until complete (or almost complete) pain resolution before resuming walking. Repeat this effort–rest cycle over 30–60 min, depending on exercise and pain tolerance. <p><i>For people who are unable or unwilling to walk at moderate–high pain intensity</i></p> <ul style="list-style-type: none"> Walk at a speed and/or grade that induces mild (2 on the claudication pain scale) claudication pain intensity. Walk and cease the exercise at the onset of the claudication pain (1 on the claudication pain scale). Rest until complete (or almost complete) pain resolution before resuming walking. Repeat this effort–rest cycle over 30–60 min, depending on exercise and pain tolerance. <p>In addition to the monitoring of the intensity of the claudication, exercise intensity^a should also be considered during the sessions.</p> <p><i>Progression</i></p> <ul style="list-style-type: none"> First, exercise training should be set at low-to-moderate intensity. Then, if well tolerated by the patient, a gradual progression to vigorous/high exercise intensity may be proposed. In general, during training programmes, the monitoring of a progressive increase in volume, intensity, and training load should be carefully considered.
Arm-ergometer	At least 3x per week		Pain-free	<p><i>Low-to-moderate</i></p> <ul style="list-style-type: none"> W_{peak}: 50%–70% HR_{max}: ≤ 76% RPE: ≤ 13 	<p><i>First weeks of training</i></p> <ul style="list-style-type: none"> One minute effort at low-to-moderate exercise intensity interspersed with 1 or 2 min rest. Repeat this effort–rest cycle four times, depending on exercise tolerance.
Cycle-ergometer	At least 3x per week		Mild to moderate	<p><i>Vigorous</i></p> <ul style="list-style-type: none"> W_{peak}: 70%–100% HR_{peak}: 77%–95% RPE: ≥ 14 	<p><i>Progression</i></p> <ul style="list-style-type: none"> Two minute effort at moderate-to-vigorous or vigorous exercise intensity interspersed with 1 or 2 min rest. Repeat this effort–rest cycle to 8–12 times, depending on exercise tolerance.

Continued

Table 4 Continued

Training modality	Training frequency	Training duration	Claudication pain intensity	Exercise intensity ^a	Example protocols
Resistance training	At least 3x per week		Mild to moderate	<p><i>Low</i></p> <p><49% 1RM RPE: 9–11</p> <p><i>Moderate</i></p> <p>50%–69% 1RM RPE: 12–13</p> <p><i>Vigorous</i></p> <p>70%–84% 1RM RPE: 14–17</p>	<p><i>First weeks of training</i></p> <ul style="list-style-type: none"> One to two sets of 12–15 repetitions (6–8 exercises) performed at low-to-moderate exercise intensity. <p><i>Progression 1</i></p> <ul style="list-style-type: none"> Two to three sets of 8–12 repetitions (6–8 exercises) performed at moderate-to-vigorous intensity. <p><i>Progression 2</i></p> <ul style="list-style-type: none"> Two to four sets of 6–8 repetitions (6–8 exercises) performed at vigorous intensity. <p><i>Example of exercises targeting the major muscle groups of the upper and lower body</i></p> <p>Leg press, knee flexion, knee extension, calf press, chest press, seated row, hip abduction, and hip extension.</p>

HR_{peak}, peak heart rate; RPE, rate of perceived exertion [Borg's scale (6, 'very very light'; 20, 'maximal effort')]; W_{peak}, peak workload; 1RM, one repetition maximum.

^aAccording to the American College of Sports Medicine guidelines for exercise testing and prescription.

Data on sex-specific differences in the efficacy of HBT are inconsistent.^{133,134} In females, the efficacy of HBT appears to be more strongly related to the individual training intensity than in males.¹³⁵ Regarding comorbidities, HBT seems to be less effective in patients with diabetes with respect to the potential increase in walking capacity.¹³⁶ In elderly patients, HBT potentially improves quality of life to a similar extent as revascularization does.¹³⁷ Considering the frequency of HBT training, 3 weekly sessions were the most commonly training strategy (range: 3 weekly sessions to daily sessions).¹²⁰ For initiation, patients should start with a duration of 20 min per session, progressively increasing the duration to 60 min per session. Home-based training can be performed outside, around a track, or in a hallway at a self-selected pace.^{51,138}


Long-term adherence to exercise therapy

In clinical practice, long-term adherence to therapy is a major problem. Participating in SET programmes may help patients to acquire awareness of the disease and learn the importance of exercise and how to practice it. Supervised exercise training programmes can be regarded as a transition phase to improve self-management and may serve as a bridge for those patients that need it to other forms of exercise approach such as community- or home-based exercise. Telemedical monitoring through step counting with pedometers or activity monitors proved to be effective,^{139,140} as did supervised structured walking exercise to improve PFWD and MWD.¹²⁰ In addition to monitoring, factors such as education, self-efficacy, goal setting, feedback, and a training plan were critical to successful outcomes.¹²⁰ This should be used more frequently in clinical practice to increase long-term adherence but needs to be demonstrated in long-term studies.

Mechanisms of response to exercise in peripheral artery disease

Exercise represents a major challenge to whole-body homeostasis provoking widespread perturbations in numerous cells, tissues, and organs that are caused by or are in response to the increased bio-energetic activity of the contracting skeletal musculature.¹⁴¹ The exercise training-induced increase in functional capacity and the concomitant amelioration of diverse maladaptive responses that ultimately reduce claudication symptoms in patients with PAD are underpinned by several inter-dependent physiological, metabolic, and mechanical mechanisms. After several months of exercise training, there is extensive remodelling of the vascular system, and although direct sampling of the vasculature in humans *in vivo* is limited, the trained musculature provides a valid proxy, being the primary tissue involved in training adaptation.¹⁴¹ The dynamic biochemical and mechanical environment around blood vessels arising from the forces provoked during skeletal muscle contractile activity (i.e. shear stress and passive stretch) and signals stimulated by the increases in muscle energetic demand (i.e. increases in AMP concentration and reduced oxygen delivery) activate several intracellular signalling pathways responsible for promoting a regulatory network governing the transcriptional control of mitochondrial biogenesis and respiratory function along with enhanced expression of pro-angiogenic factors¹⁴² (Figure 3).

Over time, this results in the initiation of capillary growth and a proliferation in the number of arterioles. Such structural remodelling is driven by a complex and often redundant sequence of events that include NO and prostaglandins. Indeed, mechanical, neural, and humoral factors, including those released from contracting skeletal muscle, have all been implicated in the remodelling response, with the vascular endothelial growth factor (VEGF) signalling pathway and downstream targets ultimately driving skeletal muscle capillary expansion.¹⁴² Muscle



activity increases VEGF in the muscle interstitium and subsequently acts on the VEGF receptors, VEGFR-1 and VEGFR-2, on the capillary endothelium, activating multiple downstream pathways via signalling intermediates such as mitogen-activated protein kinases (MAPK) and phosphatidylinositol-3-Kinase.¹⁴³ The time-course of remodelling varies and is largely a function of the blood vessel size, and while many of these adaptations are restricted to the vascular beds of the trained muscles, improved endothelial function appears to be a whole-body response to exercise training, even in individuals with PAD.

Vascular endothelial growth factor expression is partially regulated by the hypoxia-inducible factor-1 α (HIF-1 α), but recently, the peroxisome proliferator-activated receptor gamma coactivator-1 α (PGC-1 α) has emerged as an important candidate in the exercise-induced angiogenic response. PGC-1 α regulates the coordinated expression of mitochondrial proteins encoded in the nuclear and mitochondrial genomes and is rapidly induced after exercise. This protein has been called the 'master regulator' of mitochondrial biogenesis and controls various aspects of muscle oxidative phenotype while transducing and integrating physiological signals governing metabolism, differentiation, and cell growth and suppressing a broad inflammatory response.¹⁴⁴ Thus, the PGC-1 coactivators serve as a central component of the transcriptional regulatory circuitry that coordinates the energy-generating functions of the mitochondria in accordance with the metabolic demands imposed by exercise training undertaken by patients with PAD.

Exercise and revascularization

Current guidelines recommend SET programmes as an initial treatment modality for patients with IC.^{3,145} Revascularization is recommended

for patients with IC when they do not respond to initial exercise and medical therapies.¹⁴⁶ However, the role of revascularization as an initial treatment option alone or as an upstream adjunct to SET in patients with IC remains controversial.

Several trials have compared endovascular therapies with or without SET vs. SET alone as an initial treatment strategy for patients with PAD with IC and reported inconsistent results.^{147–150}

The relevant aspect of exercise training may be the reduction of the inflammatory process in patients with PAD. In a recent trial, reactive oxygen species (ROS) formation was measured using the luminol analogue L-012 for patients with IC, randomized either to HBT alone or in addition to endovascular therapy (EVT).¹⁵¹ Follow-up was performed after 3 months. Reactive oxygen species production after NAPDH oxidase 2 (NOX2) stimulation showed a significant reduction in both groups at follow-up (EVT group, $P = .002$; exercise group, $P = .019$), with a higher relative reduction in ROS in the EVT group than in the exercise group ($P = .014$).

The data regarding the benefit of SET alone or in combination with EVT or EVT alone are rare. A robust evaluation of existing data comes from a meta-analysis comparing the different treatment approaches.¹⁵² A total of 987 patients from 7 randomized control trials (constituting 9 total comparison arms) with a median follow-up duration of 12.4 months (range 10–18 months) were enrolled. Of these, 530 patients were randomized to EVT vs. SET alone and 457 patients to EVT plus SET vs. SET alone.¹⁵² For the effect of EVT alone vs. SET alone (five comparison arms), a random effects model showed no significant difference in the MWD [standardized mean difference (SMD): -0.11 (95% CI: -0.59 to 0.36); $P = .64$] on follow-up between the two groups,

neither for the PFWD, need for revascularization, or amputation. On pooled analysis, the ABI was significantly higher among participants that underwent EVT alone as compared with SET only [SMD: 0.64; 95% CI: 0.38–0.90, $P < .0001$; weighted mean difference (WMD): 0.15; 95% CI: 0.10–0.19, $P < .0001$].

On pooled analysis using random effects models, EVT plus SET (four comparison arms) was associated with significantly higher MWD on follow-up compared with SET alone (SMD: 0.79; 95% CI: 0.18–1.39, $P = .01$), as well as significantly higher ABI on follow-up compared with SET only (SMD: 0.62; 95% CI: 0.33–0.91; WMD: 0.14; 95% CI: 0.10–0.17, $P < .0001$).

The combination of EVT plus SET was also associated with a significantly lower risk of revascularization or amputation on follow-up [3.5% vs. 17.3%, odds ratio (OR): 0.19; 95% CI: 0.09–0.40, $P < .0001$]. The corresponding number needed to treat was eight patients (95% CI: 6–12). Pain-free walking distance was reported in two studies with no difference between the two groups in random effects pooled analysis.¹⁵² However, EVT alone is not associated with better outcomes than SET.^{152,153} Among patients with stable PAD and IC, compared with SET alone, endovascular revascularization in combination with SET is associated with improved outcomes.

Exercise training after surgical revascularization also improves outcomes compared with revascularization without exercise training. Although much less investigated, few publications exist on the impact of exercise on the outcome after surgical revascularization of symptomatic PAD. One small RCT compared patients after bypass surgery ($n = 14$).¹⁵⁴ Group I had standard preoperative and postoperative care, and the intervention group (Group II) had SET 4–10 weeks postoperatively. Maximal walking distance, mean increase in ABI and improvement in WIQ were significantly better in Group II. In another recent study, patients who underwent above knee femoropopliteal bypass were divided into two groups: those who continued regular exercise after bypass operation with those who discontinued exercise after surgery.¹⁵⁵ After propensity score matching, 5-year primary and secondary patency (PP: 97% vs. 61%, $P = .0041$; SP: 100% vs. 69%, $P = .0021$) and freedom from major adverse CV events (61% vs. 24%, $P = .0071$) were significantly better in patients who continued exercise. One systematic review included all RCTs with either surgical or endovascular revascularization to evaluate the evidence on the efficacy of lower limb revascularization combined with SET in patients with PAD.¹⁵⁶ Eight trials with 726 patients showed that combined therapy led to greater improvements in PFWD and MWD compared with revascularization or supervised training alone. In two out of eight studies, revascularization was surgical and in six studies it was endovascular.

Effect of exercise on health-related quality of life and cognitive function

Poor HRQoL is associated with higher rate of mortality in patients with PAD.¹⁵⁷ Randomized controlled trials have shown that exercise training vs. usual medical care in patients with PAD improves not only the perceived walking distance and speed but also the functional status as measured by specific impairment questionnaires, as the WIQ. When compared with controls, patients who complete any form of exercise training significantly improve their WIQ speed (mean difference: 9.60; 95% CI: 6.98–12.23, $P \leq .001$), WIQ distance (mean difference: 7.41; 95% CI: 4.49–10.33, $P \leq .001$), and WIQ stair climbing (mean difference: 5.07; 95% CI: 3.16–6.99, $P \leq .001$).⁸⁰ In addition, more general HRQoL evaluation scores (Short-Form Physical Component Summary) also showed significant improvement following exercise

therapy (mean difference: 1.24; 95% CI: 0.48–2.01).⁸⁰ Most of the studies showed that 3-,^{158–160} or 6/12-month^{94,102,161} exercise training improves patient's perception of physical HRQoL, with lesser effects on mental HRQoL. However, in the current literature, findings are inconsistent^{74,80,162} and other studies did not find the same effects.^{163–165} It is interesting to note that the improvement in general HRQoL scores (as SF-36) was mainly predicted by physical functional markers, such as the distance covered during a 6MWT (6MWD) and the history of stumbling.¹⁶⁶ These data indicate that greater improvements in physical function following exercise therapy are expected to have greater improvements in self-perceived HRQoL.¹⁶⁶ It has recently been showed that improvements in 6MWD following SET are predictive of augmentations in general HRQoL in patients with PAD.⁹⁶ Interestingly, changes in treadmill performance, which are less representative of functional walking,⁴⁶ were not related to improvements in HRQoL.⁹⁶

Regular physical activity is also known to improve cognitive functioning and brain health across the lifespan.¹⁶⁷ Cross-sectional and experimental studies show that greater amounts of physical activity are linked to better cognitive function in adults, with the best performances for exercise programmes that are structured, individualized, higher intensity, longer duration, and multicomponent.¹⁶⁸ These results support a dose-dependent neuroprotective relationship between physical exercise and cognitive performance. Physical exercise interventions aimed at improving brain health through neuroprotective mechanisms show promise for preserving cognitive performance.¹⁶⁸ Scientific evidence based on functional and neuroimaging approach has demonstrated that this relation could be mediated by improved brain integrity, including adaptations in cerebral blood flow, volume, and white matter integrity.¹⁶⁹

Patient education

All patients with PAD should be offered oral and written information about their disease so they can share decision-making and understand what they can do to help manage their condition. The role of exercise should be clearly explained, and patients should be supported to exercise regularly (assuming no contraindications). The impact of patient education regarding exercise is probably dependent on several factors, including the specific information that is provided, the timing and mode of delivery, and the nature of any interventions that are delivered concomitantly (e.g. SET). Patient education in the form of brief exercise advice, when delivered in isolation, confers little benefit and results in minimal improvement in individuals' walking distances.¹⁷⁰ Structured education programmes, on the other hand, may have greater potential to improve exercise behaviour and walking distances by building the knowledge and skills of patients to enable them to successfully self-manage their condition.¹⁷¹ Key programme features include a structured evidence-based curriculum that includes content on the nature of the condition and the role of exercise, delivery by trained educators, and embedded quality assurance processes.¹⁷¹

A systematic review by Abaraogu et al.¹⁷¹ identified 6 studies (1087 participants) that had investigated the effects of structured education for patients with PAD and IC. The interventions varied widely, but all included education sessions, exercise prescription, and behaviour change techniques. Four trials reported improvements in walking ability in intervention vs. control comparisons.¹⁷¹ Effects on physical activity and quality of life were mixed. Overall, the evidence was inconclusive and more rigorous trials are needed that include a clear and complete description of the education intervention. Participant feedback from three studies highlights intervention features that may be important

for improving physical activity, providing information about PAD/IC and exercise, providing encouragement and support with self-monitoring, and having group interaction while allowing space for individual discussion.¹⁷¹

Three other trials have tested exercise programmes that had an educational component in patients with PAD.^{172–174} The GOALS trial¹⁷³ randomized 194 participants either to a group-mediated cognitive behavioural intervention or an attention control group. The intervention consisted of group meetings with a facilitator once weekly for 6 months. Discussion topics included effective behaviour change methods, self-monitoring, exercising in cold weather, managing leg pain during exercise, and overcoming other obstacles to exercise adherence. At the 6-month follow-up, the intervention group achieved a 53.5 m greater increase in 6MWD compared with the control group. Next, the HONOR trial¹⁷⁴ tested the efficacy of telephone coaching combined with a wearable activity monitor and showed no improvement in 6MWD at the 9-month follow-up. Finally, the MOSAIC trial explored the effect of a physiotherapist-delivered motivational interviewing intervention in 190 patients with PAD and IC.¹⁷² A statistically significant mean difference of 16.7 m in 6MWD was observed at 3 months follow-up compared with usual care control.¹⁷² The contrasting results of these trials indicate that exercise programmes that include education are more likely to be successful if they include periodic visits to a medical centre to meet with a coach or include tailored behaviour change components.

Sex and exercise

Prevalence of PAD in women is similar to men at all ages.^{175,176} However, women are more likely to have asymptomatic PAD and less likely to report IC.¹⁷⁷ Decreased detection and subsequent intervention may then result in a higher proportion of women with severe disease and chronic limb-threatening ischaemia. Further, women who undergo revascularization tend to be older and have more severe PAD compared with men, and these factors can affect outcomes of procedures adversely.¹⁷⁸ Contradictory results exist on women with PAD and mortality rates.^{179–181} Population studies suggest a trend towards higher mortality rates in women with lower ABI.¹⁸⁰

Exercise performance has been used to suggest that women decline faster in terms of functional ability once PAD is established. However, this difference may in fact merely be due to the smaller muscles in the calves of women.¹⁸² McDermott et al.¹⁸³ showed that at 4 years of follow-up, women were more likely to become unable to walk for 6 min continuously than men, more likely to develop mobility disability, and had faster declines in walking velocity, and the distance achieved in the 6MWT was less. However, these apparent sex differences in functional decline were attenuated after additional adjustment for baseline calf muscle area and so may be attributable to smaller baseline calf muscle area in women. Interestingly poorer leg strength is associated with increased mortality in men, but not in women, with PAD.¹⁸²

The data on the efficacy of exercise rehabilitation in women with PAD compared with men are scarce. What is known, however, is that women with IC seem to have a poorer response to exercise rehabilitation, smaller changes in PFWD and MWD following 3 months of exercise than men (Δ 280 m for men vs. Δ 220 m for women; $P = .04$).¹⁸⁴ This is particularly so in those with diabetes.¹³³ Reduced blood volume expansion and slower oxygen kinetics occur in the calf musculature during exercise in women with PAD with IC.¹⁸⁵ Further, recent data showed that this poor response to exercise in women with IC and diabetes was not related to where the intervention was

performed, being impaired both in a supervised exercise class and a home exercise setting.¹³³ This poorer response to exercise was also demonstrated in the EXITPAD study, which showed that women with IC, independent of confounding factors including diabetes, benefit less from supervised exercise and have significantly lower MWD after 12 months. Higher level of metabolic syndrome present in postmenopausal women compared with similarly aged men may contribute to this.¹⁸⁴ On the contrary, it has recently been shown that multimodal SET (combining strengthening of lower limbs and Nordic walking) significantly improves walking performance (treadmill and overground) in women and men, with no difference between groups.^{98,186} Although not significant, it is interesting to note that women had greater improvements (i.e. delta) than men.⁹⁸

The clinical implication is that women with IC may respond less well to current exercise interventions and either need a greater 'dose' of exercise or another intervention separate or in combination with exercise, to obtain similar improvements in IC as that seen in men with exercise alone.

Situation in Europe

Despite of the large body of evidence highlighting benefits, SET is underused, and its availability and adherence is low.^{129–131,187–193} To note, the rate of clinicians referring a patient for SET is very low.¹²⁹ The reasons and barriers for not participating in SET programmes are lack of facilities, feeling worse, costs, time, lack of motivation, and comorbidities.^{129,131,188}

The situation with SET in Europe varies from country to country. A recent European survey showed that supervised exercise programmes exist in Austria, Belgium, Czech Republic, France, Germany, Italy, Sweden, Switzerland, and UK.¹⁹⁴ However, SET is reimbursed by the health insurance only in Austria, Belgium, France, Germany, Sweden, and Switzerland.¹⁹⁴ In the UK, SET programmes are funded by the National Health Service. In contrast, SET is not reimbursed in Czech Republic and Italy, and it even does not exist for patients with PAD in Denmark, Greece, Ireland, Poland, Serbia, Slovakia, Slovenia, or Ukraine.¹⁹⁴ Similarly, the structured home-based exercise programme is not routinely present in European countries.¹⁹⁴

Importantly, there is heterogeneity in the form of SET in most of individual countries, with existence of individual programmes or practice of each hospital or community.¹⁹⁴ They differ in respect of frequency, length, and duration of training, type of exercise, as well as by supervising professional.¹⁹⁴ Mostly, the SET is coordinated by angiologist/vascular physician, but sessions are predominantly supervised by clinical exercise physiologists or physiotherapists. Supervised exercise training for patients with PAD is sometimes offered in cardiac rehabilitation centres. Training programme duration is mostly 12 weeks or less, with session duration 30–60 min. Most often used training modalities are combination of walking and resistance training or walking training alone.¹⁹⁴

To standardize SET programmes and provision across Europe, the following steps are required: (i) a more widespread availability of SET programmes and standardized outcomes to assess their effectiveness; (ii) a more defined harmonization of SET characteristics (establish process of referral, supervision, coordination, selection of patients, and SET protocols); (iii) health insurance reimbursement of costs; and (iv) action to improve the public knowledge about the benefits of SET.¹⁹⁴

Gaps in evidence and further studies

Awareness and access to supervised exercise programmes should be a field of further studies. Additionally, there are still many areas of

insufficient or inconsistent evidence in the treatment of claudication with exercise therapy. We do not know the optimal therapy in terms of duration of the single walking session or intensity of training. We have few studies on the impact of no or low pain-based exercise, and the data on sex differences are inconsistent. The combination of walking exercise with non-walking training has not been yet established. Also, we need more evidence to better understand the potential role of wearable monitoring during exercise interventions and to evaluate on the efficacy of supportive interventions that can be used together with exercise therapy. For example, the effect of different hydration strategies used during exercise training needs more evidence. In a non-randomized study, Parodi et al.¹⁹⁵ reported mean increase in treadmill walking from 100 to 535 m in 131 patients, who were treated with hydration, determined as drinking at least 2000 mL of water during 24 h for a period of 6 months and ingesting albumin and salt (3.5 g/day).

Moreover, data on the interference of exercise training, as well as of individual training modalities, with medical treatment in patients with IC are scarce: one historic RCT suggested an augmentation of the beneficial effect of exercise training by antiplatelet therapy.¹⁹⁶ Another more recent RCT suggested an additive effect of cilostazol on top of exercise treatment on absolute claudication distance.¹⁹⁷ However, it needs to be taken into account that both studies had very small sample sizes. Therefore, larger prospective trials are needed to further elucidate the interaction between exercise training and medication in PAD.

Another area of future research should be exploration of the best modalities to transition patients from supervised exercise programmes to everyday life while maintaining the beneficial effects. Finally, we need more research on how to measure success in exercise training in an accurate and reproducible way.

Acknowledgements

The authors would like to thank Vinko Boc (Department of Vascular Diseases, University Medical Centre Ljubljana, 1000 Ljubljana, Slovenia); Tristan Mirault (Université Paris Cité, Inserm, PARCC, F-75015, Paris, France; Service de médecine vasculaire, Hôpital Européen G. Pompidou, Paris, France); Frederico Bastos Gonçalves (Department of Angiology and Vascular Surgery, Hospital de Santa Marta, Centro Hospitalar de Lisboa Central, Lisbon, Portugal); and Christian-Alexander Behrendt (Department of Vascular and Endovascular Surgery, Asklepios Clinic Wandsbek, Asklepios Medical School, Hamburg, Germany) for critically reviewing the paper.

Supplementary Data

Supplementary data are not available at *European Heart Journal* online.

Declarations

Disclosure of Interest

All authors declare no disclosure of interest for this contribution.

Data Availability

No data were generated or analysed for this manuscript.

Funding

All authors declare no funding for this contribution.

References

- Pelliccia A, Sharma S, Gati S, Back M, Borjesson M, Caselli S, et al. 2020 ESC guidelines on sports cardiology and exercise in patients with cardiovascular disease. *Eur Heart J* 2021;**42**:17–96. <https://doi.org/10.1093/eurheartj/ehaa605>
- Song P, Rudan D, Zhu Y, Fowkes FJL, Rahimi K, Fowkes FGR, et al. Global, regional, and national prevalence and risk factors for peripheral artery disease in 2015: an updated systematic review and analysis. *Lancet Glob Health* 2019;**7**:e1020–30. [https://doi.org/10.1016/S2214-109X\(19\)30255-4](https://doi.org/10.1016/S2214-109X(19)30255-4)
- Aboyans V, Ricco JB, Bartelink MEL, Björck M, Brodmann M, Cohnert T, et al. 2017 ESC guidelines on the diagnosis and treatment of peripheral arterial diseases, in collaboration with the European Society for Vascular Surgery (ESVS): document covering atherosclerotic disease of extracranial carotid and vertebral, mesenteric, renal, upper and lower extremity arteries. Endorsed by: the European Stroke Organization (ESO) The Task Force for the Diagnosis and Treatment of Peripheral Arterial Diseases of the European Society of Cardiology (ESC) and of the European Society for Vascular Surgery (ESVS). *Eur Heart J* 2018;**39**:763–816. <https://doi.org/10.1093/eurheartj/ehx095>
- Frank U, Nikol S, Belch J, Boc V, Brodmann M, Carpentier PH, et al. ESVS guideline on peripheral arterial disease. *Vasa* 2019;**48**:1–79. <https://doi.org/10.1024/0301-1526/a000834>
- McDermott MM. Lower extremity manifestations of peripheral artery disease: the pathophysiologic and functional implications of leg ischemia. *Circ Res* 2015;**116**:1540–50. <https://doi.org/10.1161/CIRCRESAHA.114.303517>
- McDermott MM, Greenland P, Liu K, Guralnik JM, Celis L, Criqui MH, et al. The ankle brachial index is associated with leg function and physical activity: the Walking and Leg Circulation Study. *Ann Intern Med* 2002;**136**:873–83. <https://doi.org/10.7326/0003-4819-136-12-200206180-00008>
- McDermott MM, Liu K, Ferrucci L, Tian L, Guralnik JM, Liao Y, et al. Decline in functional performance predicts later increased mobility loss and mortality in peripheral arterial disease. *J Am Coll Cardiol* 2011;**57**:962–70. <https://doi.org/10.1016/j.jacc.2010.09.053>
- Weitz JJ, Byrne J, Clagett GP, Farkouh ME, Porter JM, Sackett DL, et al. Diagnosis and treatment of chronic arterial insufficiency of the lower extremities: a critical review. *Circulation* 1996;**94**:3026–49. <https://doi.org/10.1161/01.CIR.94.11.3026>
- Rose GA. The diagnosis of ischaemic heart pain and intermittent claudication in field surveys. *Bull World Health Organ* 1962;**27**:645–58.
- McDermott MM, Mehta S, Greenland P. Exertional leg symptoms other than intermittent claudication are common in peripheral arterial disease. *Arch Intern Med* 1999;**159**:387–92. <https://doi.org/10.1001/archinte.159.4.387>
- Hiatt WR, Armstrong EJ, Larson CJ, Brass EP. Pathogenesis of the limb manifestations and exercise limitations in peripheral artery disease. *Circ Res* 2015;**116**:1527–39. <https://doi.org/10.1161/CIRCRESAHA.116.303566>
- Seretny M, Colvin LA. Pain management in patients with vascular disease. *Br J Anaesth* 2016;**117**:ii95–ii106. <https://doi.org/10.1093/bja/aew212>
- Tew GA, Ouedraogo N, Nicolas G, Leftheriotis G, Copeland RJ, Abraham P. Impaired somatosensation in patients with isolated proximal-without-distal exercise-related lower-limb ischemia. *Clin J Pain* 2012;**28**:404–9. <https://doi.org/10.1097/AJP.0b013e3182340c01>
- Hammad TA, Strefling JA, Zellers PR, Reed GW, Venkatachalam S, Lowry AM, et al. The effect of post-exercise ankle-brachial index on lower extremity revascularization. *JACC Cardiovasc Interv* 2015;**8**:1238–44. <https://doi.org/10.1016/j.jcin.2015.04.021>
- McDermott MM, Dayanidhi S, Kosmac K, Saini S, Slysz J, Leeuwenburgh C, et al. Walking exercise therapy effects on lower extremity skeletal muscle in peripheral artery disease. *Circ Res* 2021;**128**:1851–67. <https://doi.org/10.1161/CIRCRESAHA.121.318242>
- Sheikh MA, Bhatt DL, Li J, Lin S, Bartholomew JR. Usefulness of postexercise ankle-brachial index to predict all-cause mortality. *Am J Cardiol* 2011;**107**:778–82. <https://doi.org/10.1016/j.amjcard.2010.10.060>
- Flammer AJ, Anderson T, Celermajer DS, Creager MA, Deanfield J, Ganz P, et al. The assessment of endothelial function: from research into clinical practice. *Circulation* 2012;**126**:753–67. <https://doi.org/10.1161/CIRCULATIONAHA.112.093245>
- Gokce N, Vita JA, Bader DS, Sherman DL, Hunter LM, Holbrook M, et al. Effect of exercise on upper and lower extremity endothelial function in patients with coronary artery disease. *Am J Cardiol* 2002;**90**:124–7. [https://doi.org/10.1016/S0002-9149\(02\)02433-5](https://doi.org/10.1016/S0002-9149(02)02433-5)
- Meredith IT, Currie KE, Anderson TJ, Roddy MA, Ganz P, Creager MA. Postischemic vasodilation in human forearm is dependent on endothelium-derived nitric oxide. *Am J Physiol* 1996;**270**:H1435–40. <https://doi.org/10.1152/ajpheart.1996.270.4.H1435>
- Vita JA, Hamburg NM. Does endothelial dysfunction contribute to the clinical status of patients with peripheral arterial disease? *Can J Cardiol* 2010;**26**:45A–50A. [https://doi.org/10.1016/S0828-282X\(10\)71062-X](https://doi.org/10.1016/S0828-282X(10)71062-X)
- Robbins JL, Jones WS, Duscha BD, Allen JD, Kraus WE, Regensteiner JG, et al. Relationship between leg muscle capillary density and peak hyperemic blood flow with endurance capacity in peripheral artery disease. *J Appl Physiol* (1985) 2011;**111**:81–6. <https://doi.org/10.1152/jappphysiol.00141.2011>

22. Beckman JA, Preis O, Ridker PM, Gerhard-Herman M. Comparison of usefulness of inflammatory markers in patients with versus without peripheral arterial disease in predicting adverse cardiovascular outcomes (myocardial infarction, stroke, and death). *Am J Cardiol* 2005;**96**:1374–8. <https://doi.org/10.1016/j.amjcard.2005.07.041>
23. Tzoulaki I, Murray GD, Lee AJ, Rumley A, Lowe GD, Fowkes FG. C-reactive protein, interleukin-6, and soluble adhesion molecules as predictors of progressive peripheral atherosclerosis in the general population: Edinburgh Artery Study. *Circulation* 2005;**112**:976–83. <https://doi.org/10.1161/CIRCULATIONAHA.104.513085>
24. Vidula H, Tian L, Liu K, Criqui MH, Ferrucci L, Pearce WH, et al. Biomarkers of inflammation and thrombosis as predictors of near-term mortality in patients with peripheral arterial disease: a cohort study. *Ann Intern Med* 2008;**148**:85–93. <https://doi.org/10.7326/0003-4819-148-2-200801150-00003>
25. McDermott MM, Ferrucci L, Liu K, Guralnik JM, Tian L, Liao Y, et al. Leg symptom categories and rates of mobility decline in peripheral arterial disease. *J Am Geriatr Soc* 2010;**58**:1256–62. <https://doi.org/10.1111/j.1532-5415.2010.02941.x>
26. Gardner AW, Forrester L, Smith GV. Altered gait profile in subjects with peripheral arterial disease. *Vasc Med* 2001;**6**:31–4. <https://doi.org/10.1177/1358836X0100600106>
27. Gommans LNM, Smid AT, Scheltinga MRM, Cancrinus E, Brooijmans FAM, Meijer K, et al. Prolonged stance phase during walking in intermittent claudication. *J Vasc Surg* 2017;**66**:515–22. <https://doi.org/10.1016/j.jvs.2017.02.033>
28. McDermott MM, Liu K, Greenland P, Guralnik JM, Criqui MH, Chan C, et al. Functional decline in peripheral arterial disease: associations with the ankle brachial index and leg symptoms. *JAMA* 2004;**292**:453–61. <https://doi.org/10.1001/jama.292.4.453>
29. Gardner AW, Montgomery PS, Ritti-Dias RM, Forrester L. The effect of claudication pain on temporal and spatial gait measures during self-paced ambulation. *Vasc Med* 2010;**15**:21–6. <https://doi.org/10.1177/1358863X09106836>
30. Koutakis P, Johanning JM, Haynatzki GR, Myers SA, Stergiou N, Longo GM, et al. Abnormal joint powers before and after the onset of claudication symptoms. *J Vasc Surg* 2010;**52**:340–7. <https://doi.org/10.1016/j.jvs.2010.03.005>
31. Koutakis P, Pipinos II, Myers SA, Stergiou N, Lynch TG, Johanning JM. Joint torques and powers are reduced during ambulation for both limbs in patients with unilateral claudication. *J Vasc Surg* 2010;**51**:80–8. <https://doi.org/10.1016/j.jvs.2009.07.117>
32. Schieber MN, Hasenkamp RM, Pipinos II, Johanning JM, Stergiou N, DeSpiegelaere HK, et al. Muscle strength and control characteristics are altered by peripheral artery disease. *J Vasc Surg* 2017;**66**:178–86 e12. <https://doi.org/10.1016/j.jvs.2017.01.051>
33. Gardner AW, Montgomery PS. Impaired balance and higher prevalence of falls in subjects with intermittent claudication. *J Gerontol A Biol Sci Med Sci* 2001;**56**:M454–8. <https://doi.org/10.1093/gerona/56.7.M454>
34. Gohil RA, Mockford KA, Mazari F, Khan J, Vanicek N, Chetter IC, et al. Balance impairment, physical ability, and its link with disease severity in patients with intermittent claudication. *Ann Vasc Surg* 2013;**27**:68–74. <https://doi.org/10.1016/j.avsg.2012.05.005>
35. Chaudru S, Jehannin P, de Mullenheim PY, Klein H, Jaquinandi V, Mahe G, et al. Using wearable monitors to assess daily walking limitations induced by ischemic pain in peripheral artery disease. *Scand J Med Sci Sports* 2019;**29**:1813–26. <https://doi.org/10.1111/sms.13511>
36. McDermott MM, Guralnik JM, Tian L, Liu K, Ferrucci L, Liao Y, et al. Associations of borderline and low normal ankle-brachial index values with functional decline at 5-year follow-up: the WALCS (Walking and Leg Circulation Study). *J Am Coll Cardiol* 2009;**53**:1056–62. <https://doi.org/10.1016/j.jacc.2008.09.063>
37. Abovans V, Criqui MH, Abraham P, Allison MA, Creager MA, Diehm C, et al. Measurement and interpretation of the ankle-brachial index: a scientific statement from the American Heart Association. *Circulation* 2012;**126**:2890–909. <https://doi.org/10.1161/CIR.0b013e318276fbc6>
38. Treat-Jacobson D, McDermott MM, Beckman JA, Burt MA, Creager MA, Ehrman JK, et al. Implementation of supervised exercise therapy for patients with symptomatic peripheral artery disease: a science advisory from the American Heart Association. *Circulation* 2019;**140**:e700–e10. <https://doi.org/10.1161/CIR.0000000000000727>
39. Gardner AW, Skinner JS, Cantwell BV, Smith LK. Progressive vs single-stage treadmill tests for evaluation of claudication. *Med Sci Sports Exerc* 1991;**23**:402–8. <https://doi.org/10.1249/00005768-199104000-00003>
40. Hiatt WR, Hirsch AT, Regensteiner JG, Brass EP. Clinical trials for claudication. Assessment of exercise performance, functional status, and clinical end points. *Vascular Clinical Trialists. Circulation* 1995;**92**:614–21. <https://doi.org/10.1161/01.CIR.92.3.614>
41. Hiatt WR, Rogers RK, Brass EP. The treadmill is a better functional test than the 6-minute walk test in therapeutic trials of patients with peripheral artery disease. *Circulation* 2014;**130**:69–78. <https://doi.org/10.1161/CIRCULATIONAHA.113.007003>
42. Nicolai SP, Viechtbauer W, Kruidenier LM, Candel MJ, Prins MH, Teijink JA. Reliability of treadmill testing in peripheral arterial disease: a meta-regression analysis. *J Vasc Surg* 2009;**50**:322–9. <https://doi.org/10.1016/j.jvs.2009.01.042>
43. ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med* 2002;**166**:111–7. <https://doi.org/10.1164/ajrccm.166.1.at1102>
44. Chandra D, Kulkarni HS, Sciruba F. Learning from the learning effect in the six-minute-walk test. *Am J Respir Crit Care Med* 2012;**185**:684. <https://doi.org/10.1164/ajrccm.185.6.684>
45. Tew G, Copeland R, Le Faucheur A, Gernigon M, Nawaz S, Abraham P. Feasibility and validity of self-reported walking capacity in patients with intermittent claudication. *J Vasc Surg* 2013;**57**:1227–34. <https://doi.org/10.1016/j.jvs.2012.02.073>
46. McDermott MM, Guralnik JM, Criqui MH, Liu K, Kibbe MR, Ferrucci L. Six-minute walk is a better outcome measure than treadmill walking tests in therapeutic trials of patients with peripheral artery disease. *Circulation* 2014;**130**:61–8. <https://doi.org/10.1161/CIRCULATIONAHA.114.007002>
47. McDermott MM, Ades PA, Dyer A, Guralnik JM, Kibbe M, Criqui MH. Corridor-based functional performance measures correlate better with physical activity during daily life than treadmill measures in persons with peripheral arterial disease. *J Vasc Surg* 2008;**48**:1231–7. <https://doi.org/10.1016/j.jvs.2008.06.050>
48. McDermott MM, Guralnik JM, Tian L, Ferrucci L, Liu K, Liao Y, et al. Baseline functional performance predicts the rate of mobility loss in persons with peripheral arterial disease. *J Am Coll Cardiol* 2007;**50**:974–82. <https://doi.org/10.1016/j.jacc.2007.05.030>
49. Sandberg A, Cider A, Jivegard L, Nordanstig J, Wittboldt S, Back M. Test-retest reliability, agreement, and minimal detectable change in the 6-minute walk test in patients with intermittent claudication. *J Vasc Surg* 2020;**71**:197–203. <https://doi.org/10.1016/j.jvs.2019.02.056>
50. McDermott MM, Tian L, Criqui MH, Ferrucci L, Conte MS, Zhao L, et al. Meaningful change in 6-minute walk in people with peripheral artery disease. *J Vasc Surg* 2021;**73**:267–76 e1. <https://doi.org/10.1016/j.jvs.2020.03.052>
51. Gardner AW, Montgomery PS, Wang M. Minimal clinically important differences in treadmill, 6-minute walk, and patient-based outcomes following supervised and home-based exercise in peripheral artery disease. *Vasc Med* 2018;**23**:349–57. <https://doi.org/10.1177/1358863X18762599>
52. de Mullenheim PY, Chaudru S, Mahe G, Prioux J, Le Faucheur A. Clinical interest of ambulatory assessment of physical activity and walking capacity in peripheral artery disease. *Scand J Med Sci Sports* 2016;**26**:716–30. <https://doi.org/10.1111/sms.12512>
53. Abraham P, Noury-Desvaux B, Gernigon M, Mahe G, Sauvaget T, Leftheriotis G, et al. The inter- and intra-unit variability of a low-cost GPS data logger/receiver to study human outdoor walking in view of health and clinical studies. *PLoS One* 2012;**7**:e31338. <https://doi.org/10.1371/journal.pone.0031338>
54. Taoum A, Chaudru S, DE Mullenheim PY, Congnard F, Emily M, Noury-Desvaux B, et al. Comparison of activity monitors accuracy in assessing intermittent outdoor walking. *Med Sci Sports Exerc* 2021;**53**:1303–14. <https://doi.org/10.1249/MSS.0000000000002587>
55. Hochsmann C, Knaier R, Eymann J, Hintermann J, Infanger D, Schmidt-Trucksass A. Validity of activity trackers, smartphones, and phone applications to measure steps in various walking conditions. *Scand J Med Sci Sports* 2018;**28**:1818–27. <https://doi.org/10.1111/sms.13074>
56. Le Faucheur A, Abraham P, Jaquinandi V, Bouye P, Saumet JL, Noury-Desvaux B. Measurement of walking distance and speed in patients with peripheral arterial disease: a novel method using a global positioning system. *Circulation* 2008;**117**:897–904. <https://doi.org/10.1161/CIRCULATIONAHA.107.725994>
57. McDermott MM, Tian L, Ferrucci L, Liu K, Guralnik JM, Liao Y, et al. Associations between lower extremity ischemia, upper and lower extremity strength, and functional impairment with peripheral arterial disease. *J Am Geriatr Soc* 2008;**56**:724–9. <https://doi.org/10.1111/j.1532-5415.2008.01633.x>
58. Pizzimenti M, Meyer A, Charles AL, Giannini M, Chakfe N, Lejay A, et al. Sarcopenia and peripheral arterial disease: a systematic review. *J Cachexia Sarcopenia Muscle* 2020;**11**:866–86. <https://doi.org/10.1002/jcsm.12587>
59. Ritti-Dias RM, Basyches M, Camara L, Puech-Leao P, Battistella L, Wolosker N. Test-retest reliability of isokinetic strength and endurance tests in patients with intermittent claudication. *Vasc Med* 2010;**15**:275–8. <https://doi.org/10.1177/1358863X10371415>
60. Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol* 1994;**49**:M85–94. <https://doi.org/10.1093/geronj/49.2.M85>
61. Lanzi S, Pousaz A, Calanca L, Mazzolai L. Sit to stand muscle power is related to functional performance at baseline and after supervised exercise training in patients with lower extremity peripheral artery disease. *Eur J Vasc Endovasc Surg* 2023;**65**:521–7. <https://doi.org/10.1016/j.ejvs.2022.12.029>
62. Arndt H, Nordanstig J, Bertges DJ, Budtz-Lilly J, Venermo M, Espada CL, et al. A Delphi consensus on patient reported outcomes for registries and trials including patients with intermittent claudication: recommendations and reporting standard. *Eur J Vasc Endovasc Surg* 2022;**64**:526–33. <https://doi.org/10.1016/j.ejvs.2022.08.011>
63. Raja A, Spertus J, Yeh RW, Secemsky EA. Assessing health-related quality of life among patients with peripheral artery disease: a review of the literature and focus on patient-reported outcome measures. *Vasc Med* 2021;**26**:317–25. <https://doi.org/10.1177/1358863X20977016>
64. Treat-Jacobson D, McDermott MM, Bronas UG, Campia U, Collins TC, Criqui MH, et al. Optimal exercise programs for patients with peripheral artery disease: a scientific statement from the American Heart Association. *Circulation* 2019;**139**:e10–33. <https://doi.org/10.1161/CIR.0000000000000623>
65. Regensteiner JG, Hiatt WR, Coll JR, Criqui MH, Treat-Jacobson D, McDermott MM, et al. The impact of peripheral arterial disease on health-related quality of life in the

- Peripheral Arterial Disease Awareness, Risk, and Treatment: New Resources for Survival (PARTNERS) program. *Vasc Med* 2008;**13**:15–24. <https://doi.org/10.1177/1358863X07084911>
66. Kim M, Kim Y, Ryu GW, Choi M. Functional status and health-related quality of life in patients with peripheral artery disease: a cross-sectional study. *Int J Environ Res Public Health* 2021;**18**:10941. <https://doi.org/10.3390/ijerph182010941>
 67. Franco OH, de Laet C, Peeters A, Jonker J, Mackenbach J, Nusselder W. Effects of physical activity on life expectancy with cardiovascular disease. *Arch Intern Med* 2005;**165**:2355–60. <https://doi.org/10.1001/archinte.165.20.2355>
 68. Dostalova R, Stillman C, Erickson KI, Slepicka P, Mudrak J. The relationship between physical activity, self-perceived health, and cognitive function in older adults. *Brain Sci* 2021;**11**:492. <https://doi.org/10.3390/brainsci11040492>
 69. Frank U, Nikol S, Belch J. 5 Conservative treatment for PAD—risk factor management. *Vasa* 2019;**48**:1–12. <https://doi.org/10.1024/0301-1526/a000835>
 70. Harwood A, Pymmer S, Ingle L, Doherty P, Chetter I, Parmenter B, et al. Exercise training for intermittent claudication: a narrative review and summary of guidelines for practitioners. *BMJ Open Sport & Exercise Medicine* 2020;**6**:e000897. <https://doi.org/10.1136/bmjsem-2020-000897>
 71. Gommans LN, Fokkenrood HJ, van Dalen HC, Scheltinga MR, Teijink JA, Peters RJ. Safety of supervised exercise therapy in patients with intermittent claudication. *J Vasc Surg* 2015;**61**:512–8 e2. <https://doi.org/10.1016/j.jvs.2014.08.070>
 72. Bronas UG, Regensteiner JG. Connecting the past to the present: a historical review of exercise training for peripheral artery disease. *Vasc Med* 2022;**27**:174–85. <https://doi.org/10.1177/1358863X211073620>
 73. Penin-Grandes S, Lopez-Ortiz S, Maroto-Izquierdo S, Menendez H, Pinto-Fraga J, Martin-Hernandez J, et al. Winners do what they fear: exercise and peripheral arterial disease—an umbrella review. *Eur J Prev Cardiol* 2023;zwad261. <https://doi.org/10.1093/eurjpc/zwad261>
 74. Lane R, Harwood A, Watson L, Leng GC. Exercise for intermittent claudication. *Cochrane Database Syst Rev* 2017;**12**:CD000990. <https://doi.org/10.1002/14651858.CD000990.pub4>
 75. Fakhry F, van de Luijngaarden KM, Bax L, den Hoed PT, Hunink MG, Rouwet EV, et al. Supervised walking therapy in patients with intermittent claudication. *J Vasc Surg* 2012;**56**:1132–42. <https://doi.org/10.1016/j.jvs.2012.04.046>
 76. Lanzi S, Boichat J, Calanca L, Aubertin P, Malatesta D, Mazzolai L. Gait changes after supervised exercise training in patients with symptomatic lower extremity peripheral artery disease. *Vasc Med* 2021;**26**:259–66. <https://doi.org/10.1177/1358863X20984831>
 77. Lanzi S, Boichat J, Calanca L, Mazzolai L, Malatesta D. Supervised exercise training improves 6 min walking distance and modifies gait pattern during pain-free walking condition in patients with symptomatic lower extremity peripheral artery disease. *Sensors (Basel)* 2021;**21**:7989. <https://doi.org/10.3390/s21237989>
 78. Fassora M, Calanca L, Jaques C, Mazzolai L, Kayser B, Lanzi S. Intensity-dependent effects of exercise therapy on walking performance and aerobic fitness in symptomatic patients with lower-extremity peripheral artery disease: a systematic review and meta-analysis. *Vasc Med* 2022;**27**:158–70. <https://doi.org/10.1177/1358863X211034577>
 79. Lanzi S, Pousaz A, Calanca L, Mazzolai L. Time-course evolution of functional performance during a 3-month supervised exercise training program in patients with symptomatic peripheral artery disease. *Vasc Med* 2023;**28**:404–11. <https://doi.org/10.1177/1358863X231191908>
 80. Parmenter BJ, Dieberg G, Phipps G, Smart NA. Exercise training for health-related quality of life in peripheral artery disease: a systematic review and meta-analysis. *Vasc Med* 2015;**20**:30–40. <https://doi.org/10.1177/1358863X14559092>
 81. Parmenter BJ, Dieberg G, Smart NA. Exercise training for management of peripheral arterial disease: a systematic review and meta-analysis. *Sports Med* 2015;**45**:231–44. <https://doi.org/10.1007/s40279-014-0261-z>
 82. Schieber MN, Pipinos II, Johanning JM, Casale GP, Williams MA, DeSpiegelaere HK, et al. Supervised walking exercise therapy improves gait biomechanics in patients with peripheral artery disease. *J Vasc Surg* 2019;**71**:575–83. <https://doi.org/10.1016/j.jvs.2019.05.044>
 83. Siercke M, Jorgensen LP, Missel M, Thygesen LC, Moller SP, Sillesen H, et al. Cardiovascular rehabilitation increases walking distance in patients with intermittent claudication. Results of the CIPIC Rehab Study: a randomised controlled trial. *Eur J Vasc Endovasc Surg* 2021;**62**:768–76. <https://doi.org/10.1016/j.ejvs.2021.04.004>
 84. Brown T, Forster RB, Cleathis M, Mikhailidis DP, Stansby G, Stewart M. Cilostazol for intermittent claudication. *Cochrane Database Syst Rev* 2021;**6**:CD003748. <https://doi.org/10.1002/14651858.CD003748.pub5>
 85. De Haro J, Bleda S, Varela C, Esparza L, Acin F; Bosentan Population-Based Randomized Trial for Clinical and Endothelial Function Assessment on Endothelin Antagonist Therapy in Patients With Intermittent Claudication CLAU Investigators. Effect of bosentan on claudication distance and endothelium-dependent vasodilation in Hispanic patients with peripheral arterial disease. *Am J Cardiol* 2016;**117**:295–301. <https://doi.org/10.1016/j.amjcard.2015.10.032>
 86. Omarjee L, Le Pabic E, Custaud MA, Fontaine C, Locher C, Renault A, et al. Effects of sildenafil on maximum walking time in patients with arterial claudication: the ARTERIOFIL study. *Vascul Pharmacol* 2019;**118–119**:106563. <https://doi.org/10.1016/j.vph.2019.05.003>
 87. Suzuki J, Shimamura M, Suda H, Wakayama K, Kumagai H, Ikeda Y, et al. Current therapies and investigational drugs for peripheral arterial disease. *Hypertens Res* 2016;**39**:183–91. <https://doi.org/10.1038/hr.2015.134>
 88. Broderick C, Forster R, Abdel-Hadi M, Salhiyyah K. Pentoxifylline for intermittent claudication. *Cochrane Database Syst Rev* 2020;**10**:CD005262. <https://doi.org/10.1002/14651858.CD005262.pub4>
 89. Jansen SC, Abaraogu UO, Lauret GJ, Fakhry F, Fokkenrood HJ, Teijink JA. Modes of exercise training for intermittent claudication. *Cochrane Database Syst Rev* 2020;**8**:CD009638. <https://doi.org/10.1002/14651858.CD009638.pub3>
 90. Parmenter BJ, Raymond J, Dinnen P, Lusby RJ, Fiatarone Singh MA. High-intensity progressive resistance training improves flat-ground walking in older adults with symptomatic peripheral arterial disease. *J Am Geriatr Soc* 2013;**61**:1964–70. <https://doi.org/10.1111/jgs.12500>
 91. Ritti-Dias RM, Wolosker N, de Moraes Forjaz CL, Carvalho CR, Cucato GG, Leao PP, et al. Strength training increases walking tolerance in intermittent claudication patients: randomized trial. *J Vasc Surg* 2010;**51**:89–95. <https://doi.org/10.1016/j.jvs.2009.07.118>
 92. Sanderson B, Askew C, Stewart I, Walker P, Gibbs H, Green S. Short-term effects of cycle and treadmill training on exercise tolerance in peripheral arterial disease. *J Vasc Surg* 2006;**44**:119–27. <https://doi.org/10.1016/j.jvs.2006.03.037>
 93. Collins EG, Edwin Langbein W, Orebaugh C, Bammert C, Hanson K, Reda D, et al. Polestriding exercise and vitamin E for management of peripheral vascular disease. *Med Sci Sports Exerc* 2003;**35**:384–93. <https://doi.org/10.1249/01.MSS.0000053658.82687.FF>
 94. Collins EG, Langbein WE, Orebaugh C, Bammert C, Hanson K, Reda D, et al. Cardiovascular training effect associated with polestriding exercise in patients with peripheral arterial disease. *J Cardiovasc Nurs* 2005;**20**:177–85. <https://doi.org/10.1097/00005082-200505000-00009>
 95. Calanca L, Lanzi S, Ney B, Berchtold A, Mazzolai L. Multimodal supervised exercise significantly improves walking performances without changing hemodynamic parameters in patients with symptomatic lower extremity peripheral artery disease. *Vasc Endovascular Surg* 2020;**54**:605–11. <https://doi.org/10.1177/1538574420940090>
 96. Lanzi S, Calanca L, Berchtold A, Mazzolai L. Improvement in 6-minute walking distance after supervised exercise training is related to changes in quality of life in patients with lower extremity peripheral artery disease. *J Clin Med* 2021;**10**:3330. <https://doi.org/10.3390/jcm10153330>
 97. Lanzi S, Calanca L, Borgeat Kaeser A, Mazzolai L. Walking performances and muscle oxygen desaturation are increased after supervised exercise training in Takayasu arteritis: a case report and a review of the literature. *Eur Heart J Case Rep* 2018;**2**:yty123. <https://doi.org/10.1093/ehjcr/tyt123>
 98. Ney B, Lanzi S, Calanca L, Mazzolai L. Multimodal supervised exercise training is effective in improving long term walking performance in patients with symptomatic lower extremity peripheral artery disease. *J Clin Med* 2021;**10**:2057. <https://doi.org/10.3390/jcm10102057>
 99. Tew G, Nawaz S, Zwierska I, Saxton JM. Limb-specific and cross-transfer effects of arm-crank exercise training in patients with symptomatic peripheral arterial disease. *Clin Sci (Lond)* 2009;**117**:405–13. <https://doi.org/10.1042/CS20080688>
 100. Treat-Jacobson D, Bronas UG, Leon AS. Efficacy of arm-ergometry versus treadmill exercise training to improve walking distance in patients with claudication. *Vasc Med* 2009;**14**:203–13. <https://doi.org/10.1177/1358863X08101858>
 101. Parmenter BJ, Raymond J, Dinnen P, Singh MA. A systematic review of randomized controlled trials: walking versus alternative exercise prescription as treatment for intermittent claudication. *Atherosclerosis* 2011;**218**:1–12. <https://doi.org/10.1016/j.atherosclerosis.2011.04.024>
 102. McDermott MM, Ades P, Guralnik JM, Dyer A, Ferrucci L, Liu K, et al. Treadmill exercise and resistance training in patients with peripheral arterial disease with and without intermittent claudication: a randomized controlled trial. *JAMA* 2009;**301**:165–74. <https://doi.org/10.1001/jama.2008.962>
 103. Bulmer AC, Coombes JS. Optimising exercise training in peripheral arterial disease. *Sports Med* 2004;**34**:983–1003. <https://doi.org/10.2165/00007256-200434140-00004>
 104. Gardner AVW, Poehlman ET. Exercise rehabilitation programs for the treatment of claudication pain. A meta-analysis. *JAMA* 1995;**274**:975–80. <https://doi.org/10.1001/jama.1995.03530120067043>
 105. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 1982;**14**:377–81. <https://doi.org/10.1249/00005768-198205000-00012>
 106. NICE. The National Institute of Health and Care Excellence. Peripheral arterial disease: diagnosis and management clinical guideline [CG147]. 2012.
 107. Au TB, Golledge J, Walker PJ, Haigh K, Nelson M. Peripheral arterial disease—diagnosis and management in general practice. *Aust Fam Physician* 2013;**42**:397–400.
 108. Hammond MM, Spring B, Rejeski VJ, Sufit R, Criqui MH, Tian L, et al. Effects of walking exercise at a pace with versus without ischemic leg symptoms on functional performance measures in people with lower extremity peripheral artery disease: the LITE randomized clinical trial. *J Am Heart Assoc* 2022;**11**:e025063. <https://doi.org/10.1161/JAHA.121.025063>
 109. McDermott MM, Spring B, Tian L, Treat-Jacobson D, Ferrucci L, Lloyd-Jones D, et al. Effect of low-intensity vs high-intensity home-based walking exercise on walk distance

- in patients with peripheral artery disease: the LITE randomized clinical trial. *JAMA* 2021; **325**:1266–76. <https://doi.org/10.1001/jama.2021.2536>
110. Mika P, Konik A, Januszek R, Petriczek T, Mika A, Nowobilski R, et al. Comparison of two treadmill training programs on walking ability and endothelial function in intermittent claudication. *Int J Cardiol* 2013; **168**:838–42. <https://doi.org/10.1016/j.ijcard.2012.10.003>
 111. Novakovic M, Krevle B, Rajkovic U, Vizintin Cuderman T, Jansa Trontelj K, Fras Z, et al. Moderate-pain versus pain-free exercise, walking capacity, and cardiovascular health in patients with peripheral artery disease. *J Vasc Surg* 2019; **70**:148–56. <https://doi.org/10.1016/j.jvs.2018.10.109>
 112. Perks J, Zaccardi F, Paterson C, Houghton JSM, Nickinson ATO, Pepper CJ, et al. Effect of high-pain versus low-pain structured exercise on walking ability in people with intermittent claudication: meta-analysis. *Br J Surg* 2022; **109**:686–94. <https://doi.org/10.1093/bjs/znac134>
 113. Seed SA, Harwood AE, Sinclair J, Pymmer S, Caldwell E, Ingle L, et al. A systematic review of exercise prescription in patients with intermittent claudication: does pain matter? *Ann Vasc Surg* 2021; **77**:315–23. <https://doi.org/10.1016/j.avsg.2021.06.025>
 114. Birkett ST, Sinclair J, Seed SA, Pymmer S, Caldwell E, Ingle L, et al. Effects of exercise prescribed at different levels of claudication pain on walking performance in patients with intermittent claudication: a protocol for a randomised controlled trial. *Ther Adv Cardiovasc Dis* 2022; **16**:17539447221108817. <https://doi.org/10.1177/17539447221108817>
 115. Lanzi S, Mazzolai L. Commentary to Seed, et al. 'What is the correct level of claudication pain to prescribe? Universal inconsistency within guidelines, a painful issue. *Vascular* 2023; **17085381231160931**. <https://doi.org/10.1177/17085381231160931>
 116. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011; **43**:1334–59. <https://doi.org/10.1249/MSS.0b013e318213fefb>
 117. Hansen D, Abreu A, Ambrosetti M, Cornelissen V, Gevaert A, Kemps H, et al. Exercise intensity assessment and prescription in cardiovascular rehabilitation and beyond: why and how: a position statement from the Secondary Prevention and Rehabilitation Section of the European Association of Preventive Cardiology. *Eur J Prev Cardiol* 2022; **29**:230–45. <https://doi.org/10.1093/eurjpc/zwab007>
 118. Wood TM, Maddalozzo GF, Harter RA. Accuracy of seven equations for predicting 1-RM performance of apparently healthy, sedentary older adults. *Meas Phys Educ Exerc Sci* 2002; **6**:67–94. https://doi.org/10.1207/S15327841MPEE0602_1
 119. Parmenter BJ, Mavros Y, Ritti Dias R, King S, Fiatarone Singh M. Resistance training as a treatment for older persons with peripheral artery disease: a systematic review and meta-analysis. *Br J Sports Med* 2019; **54**:452–61. <https://doi.org/10.1136/bjsports-2018-100205>
 120. Pymmer S, Ibeggazene S, Palmer J, Tew GA, Ingle L, Smith GE, et al. An updated systematic review and meta-analysis of home-based exercise programs for individuals with intermittent claudication. *J Vasc Surg* 2021; **74**:2076–85 e20. <https://doi.org/10.1016/j.jvs.2021.03.063>
 121. van den Houten MML, Hageman D, Gommans LNM, Kleijnen J, Scheltinga MRM, Teijink JAW. The effect of supervised exercise, home based exercise and endovascular revascularisation on physical activity in patients with intermittent claudication: a network meta-analysis. *Eur J Vasc Endovasc Surg* 2019; **58**:383–92. <https://doi.org/10.1016/j.ejvs.2018.12.023>
 122. Rejeski WJ, Spring B, Domanchuk K, Tao H, Tian L, Zhao L, et al. A group-mediated, home-based physical activity intervention for patients with peripheral artery disease: effects on social and psychological function. *J Transl Med* 2014; **12**:29. <https://doi.org/10.1186/1479-5876-12-29>
 123. Fakhry F, Spronk S, de Ridder M, den Hoed PT, Hunink MG. Long-term effects of structured home-based exercise program on functional capacity and quality of life in patients with intermittent claudication. *Arch Phys Med Rehabil* 2011; **92**:1066–73. <https://doi.org/10.1016/j.apmr.2011.02.007>
 124. McDermott MM, Guralnik JM, Criqui MH, Ferrucci L, Zhao L, Liu K, et al. Home-based walking exercise in peripheral artery disease: 12-month follow-up of the GOALS randomized trial. *J Am Heart Assoc* 2014; **3**:e000711. <https://doi.org/10.1161/JAHA.113.000711>
 125. Waddell A, Seed S, Broom DR, McGregor G, Birkett ST, Harwood AE. Safety of home-based exercise for people with intermittent claudication: a systematic review. *Vasc Med* 2022; **27**:186–92. <https://doi.org/10.1177/1358863X211060388>
 126. Lamberti N, Lopez-Soto PJ, Guerzoni F, Napoli N, Gasbarro V, Zamboni P, et al. Changes in exercise capacity and risk of all-cause mortality in patients with peripheral artery disease: a 10-year retrospective cohort study. *Intern Emerg Med* 2020; **15**:289–98. <https://doi.org/10.1007/s11739-019-02176-3>
 127. Hageman D, Fokkenrood HJ, Gommans LN, van den Houten MM, Teijink JA. Supervised exercise therapy versus home-based exercise therapy versus walking advice for intermittent claudication. *Cochrane Database Syst Rev* 2018; **4**:CD005263. <https://doi.org/10.1002/14651858.CD005263.pub4>
 128. Al-Jundi W, Madbak K, Beard JD, Nawaz S, Tew GA. Systematic review of home-based exercise programmes for individuals with intermittent claudication. *Eur J Vasc Endovasc Surg* 2013; **46**:690–706. <https://doi.org/10.1016/j.ejvs.2013.09.004>
 129. Dua A, Gologorsky R, Savage D, Rens N, Gandhi N, Brooke B, et al. National assessment of availability, awareness, and utilization of supervised exercise therapy for peripheral artery disease patients with intermittent claudication. *J Vasc Surg* 2020; **71**:1702–7. <https://doi.org/10.1016/j.jvs.2019.08.238>
 130. Haque A. Few UK vascular centres offer a fully NICE-compliant supervised exercise programme: a national audit. *Ann R Coll Surg Engl* 2022; **104**:130–7. <https://doi.org/10.1308/rcsann.2021.0126>
 131. Harwood AE, Pymmer S, Ibeggazene S, Ingle L, Caldwell E, Birkett ST. Provision of exercise services in patients with peripheral artery disease in the United Kingdom. *Vascular* 2022; **30**:874–81. <https://doi.org/10.1177/17085381211035259>
 132. Makris GC, Lattimer CR, Lavid A, Geroulakos G. Availability of supervised exercise programs and the role of structured home-based exercise in peripheral arterial disease. *Eur J Vasc Endovasc Surg* 2012; **44**:569–75; discussion 76. <https://doi.org/10.1016/j.ejvs.2012.09.009>
 133. Gardner AWW, Parker DE, Montgomery PS, Blevins SM. Diabetic women are poor responders to exercise rehabilitation in the treatment of claudication. *J Vasc Surg* 2014; **59**:1036–43. <https://doi.org/10.1016/j.jvs.2013.10.058>
 134. Manfredini R, Lamberti N, Manfredini F, Straudi S, Fabbian F, Rodriguez Borrego MA, et al. Gender differences in outcomes following a pain-free, home-based exercise program for claudication. *J Womens Health (Larchmt)* 2019; **28**:1313–21. <https://doi.org/10.1089/jwh.2018.7113>
 135. Gardner AWW, Parker DE, Montgomery PS. Sex-specific predictors of improved walking with step-monitored, home-based exercise in peripheral artery disease. *Vasc Med* 2015; **20**:424–31. <https://doi.org/10.1177/1358863X15596237>
 136. Collins TC, Lunos S, Carlson T, Henderson K, Lightbourne M, Nelson B, et al. Effects of a home-based walking intervention on mobility and quality of life in people with diabetes and peripheral arterial disease: a randomized controlled trial. *Diabetes Care* 2011; **34**:2174–9. <https://doi.org/10.2337/dc10-2399>
 137. Lamberti N, Malagoni AM, Ficarra V, Basaglia N, Manfredini R, Zamboni P, et al. Structured home-based exercise versus invasive treatment: a mission impossible? A pilot randomized study in elderly patients with intermittent claudication. *Angiology* 2016; **67**:772–80. <https://doi.org/10.1177/0003319715618481>
 138. McDermott MM, Polonsky TS. Home-based exercise: a therapeutic option for peripheral artery disease. *Circulation* 2016; **134**:1127–9. <https://doi.org/10.1161/CIRCULATIONAHA.116.023691>
 139. Chan C, Sounderajah V, Normahani P, Acharya A, Markar SR, Darzi A, et al. Wearable activity monitors in home based exercise therapy for patients with intermittent claudication: a systematic review. *Eur J Vasc Endovasc Surg* 2021; **61**:676–87. <https://doi.org/10.1016/j.ejvs.2020.11.044>
 140. Kim M, Kim C, Kim E, Choi M. Effectiveness of mobile health-based exercise interventions for patients with peripheral artery disease: systematic review and meta-analysis. *JMIR Mhealth Uhealth* 2021; **9**:e24080. <https://doi.org/10.2196/24080>
 141. Hawley JA, Hargreaves M, Joyner MJ, Zierath JR. Integrative biology of exercise. *Cell* 2014; **159**:738–49. <https://doi.org/10.1016/j.cell.2014.10.029>
 142. Hoier B, Hellsten Y. Exercise-induced capillary growth in human skeletal muscle and the dynamics of VEGF. *Microcirculation* 2014; **21**:301–14. <https://doi.org/10.1111/micc.12117>
 143. Egginton S. Invited review: activity-induced angiogenesis. *Pflugers Arch* 2009; **457**:963–77. <https://doi.org/10.1007/s00424-008-0563-9>
 144. Handschin C, Spiegelman BM. The role of exercise and PGC1 α in inflammation and chronic disease. *Nature* 2008; **454**:463–9. <https://doi.org/10.1038/nature07206>
 145. Gerhard-Herman MD, Gornik HL, Barrett C, Barshes NR, Corriere MA, Drachman DE, et al. 2016 AHA/ACC guideline on the management of patients with lower extremity peripheral artery disease: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2017; **69**:1465–508. <https://doi.org/10.1016/j.jacc.2016.11.008>
 146. Heiss C, Olinic DM, Belch JFF, Brodmann M, Mazzolai L, Stanek A, et al. Management of chronic peripheral artery disease patients with indication for endovascular revascularization. *Vasa* 2022; **51**:121–37. <https://doi.org/10.1024/0301-1526/a000998>
 147. Anderson JL, Halperin JL, Albert NM, Bozkurt B, Brindis RG, Curtis LH, et al. Management of patients with peripheral artery disease (compilation of 2005 and 2011 ACCF/AHA guideline recommendations): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation* 2013; **127**:1425–43. <https://doi.org/10.1161/CIR.0b013e31828b82aa>
 148. Koelemay MJW, van Reijen NS, van Dieren S, Frans FA, Vermeulen EJJ, Buscher H, et al. Editor's choice—randomised clinical trial of supervised exercise therapy vs. endovascular revascularisation for intermittent claudication caused by iliac artery obstruction: the SUPER study. *Eur J Vasc Endovasc Surg* 2022; **63**:421–9. <https://doi.org/10.1016/j.ejvs.2021.09.042>
 149. Murphy TP, Cutlip DE, Regensteiner JG, Mohler ER, Cohen DJ, Reynolds MR, et al. Supervised exercise versus primary stenting for claudication resulting from aortoiliac

- peripheral artery disease: six-month outcomes from the Claudication: Exercise versus Endoluminal Revascularization (CLEVER) study. *Circulation* 2012;**125**:130–9. <https://doi.org/10.1161/CIRCULATIONAHA.111.075770>
150. Murphy TP, Cutlip DE, Regensteiner JG, Mohler ER III, Cohen DJ, Reynolds MR, et al. Supervised exercise, stent revascularization, or medical therapy for claudication due to aortoiliac peripheral artery disease: the CLEVER study. *J Am Coll Cardiol* 2015;**65**:999–1009. <https://doi.org/10.1016/j.jacc.2014.12.043>
 151. Koppe-Schmeisser F, Schwaderlapp M, Schmeisser J, Dopheide JF, Munzel T, Daiber A, et al. Influence of peripheral transluminal angioplasty alongside exercise training on oxidative stress and inflammation in patients with peripheral arterial disease. *J Clin Med* 2021;**10**:5851. <https://doi.org/10.3390/jcm10245851>
 152. Pandey A, Banerjee S, Ngo C, Mody P, Marso SP, Brilakis ES, et al. Comparative efficacy of endovascular revascularization versus supervised exercise training in patients with intermittent claudication: meta-analysis of randomized controlled trials. *JACC Cardiovasc Interv* 2017;**10**:712–24. <https://doi.org/10.1016/j.jcin.2017.01.027>
 153. Fakhry F, Fokkenrood HJ, Spronk S, Teijink JA, Rouwet EV, Hunink MGM. Endovascular revascularisation versus conservative management for intermittent claudication. *Cochrane Database Syst Rev* 2018;**3**:CD010512. <https://doi.org/10.1002/14651858.CD010512.pub2>
 154. Badger SA, Soong CV, O'Donnell ME, Boreham CA, McGuigan KE. Benefits of a supervised exercise program after lower limb bypass surgery. *Vasc Endovascular Surg* 2007;**41**:27–32. <https://doi.org/10.1177/1538574406296209>
 155. Kobayashi T, Hamamoto M, Okazaki T, Honma T, Iba K, Takakuwa T, et al. Effectiveness of continuous unsupervised exercise therapy after above-knee femoropopliteal bypass. *Vascular* 2021;**29**:387–95. <https://doi.org/10.1177/1708538120957488>
 156. Meneses AL, Ritti-Dias RM, Parmenter B, Gollgede J, Askew CD. Combined lower limb revascularisation and supervised exercise training for patients with peripheral arterial disease: a systematic review of randomised controlled trials. *Sports Med* 2017;**47**:987–1002. <https://doi.org/10.1007/s40279-016-0635-5>
 157. Issa SM, Hoeks SE, Scholte op Reimer VVJ, Van Gestel YR, Lenzen MJ, Verhagen HJ, et al. Health-related quality of life predicts long-term survival in patients with peripheral artery disease. *Vasc Med* 2010;**15**:163–9. <https://doi.org/10.1177/1358863X10364208>
 158. Gardner AW, Parker DE, Montgomery PS, Scott KJ, Blevins SM. Efficacy of quantified home-based exercise and supervised exercise in patients with intermittent claudication: a randomized controlled trial. *Circulation* 2011;**123**:491–8. <https://doi.org/10.1161/CIRCULATIONAHA.110.963066>
 159. Patterson RB, Pinto B, Marcus B, Colucci A, Braun T, Roberts M. Value of a supervised exercise program for the therapy of arterial claudication. *J Vasc Surg* 1997;**25**:312–8. [https://doi.org/10.1016/S0741-5214\(97\)70352-5](https://doi.org/10.1016/S0741-5214(97)70352-5)
 160. Tsai JC, Chan P, Wang CH, Jeng C, Hsieh MH, Kao PF, et al. The effects of exercise training on walking function and perception of health status in elderly patients with peripheral arterial occlusive disease. *J Intern Med* 2002;**252**:448–55. <https://doi.org/10.1046/j.1365-2796.2002.01055.x>
 161. Nicolai SP, Teijink JA, Prins MH. Exercise Therapy in Peripheral Arterial Disease Study G. Multicenter randomized clinical trial of supervised exercise therapy with or without feedback versus walking advice for intermittent claudication. *J Vasc Surg* 2010;**52**:348–55. <https://doi.org/10.1016/j.jvs.2010.02.022>
 162. Guidon M, McGee H. Exercise-based interventions and health-related quality of life in intermittent claudication: a 20-year (1989–2008) review. *Eur J Cardiovasc Prev Rehabil* 2010;**17**:140–54. <https://doi.org/10.1097/HJR.0b013e3283377f08>
 163. Guidon M, McGee H. One-year effect of a supervised exercise programme on functional capacity and quality of life in peripheral arterial disease. *Disabil Rehabil* 2013;**35**:397–404. <https://doi.org/10.3109/09638288.2012.694963>
 164. Kakkos SK, Geroulakos G, Nicolaidis AN. Improvement of the walking ability in intermittent claudication due to superficial femoral artery occlusion with supervised exercise and pneumatic foot and calf compression: a randomised controlled trial. *Eur J Vasc Endovasc Surg* 2005;**30**:164–75. <https://doi.org/10.1016/j.ejvs.2005.03.011>
 165. Savage P, Ricci MA, Lynn M, Gardner A, Knight S, Brochu M, et al. Effects of home versus supervised exercise for patients with intermittent claudication. *J Cardiopulm Rehabil* 2001;**21**:152–7. <https://doi.org/10.1097/00008483-200105000-00006>
 166. Gardner AW, Montgomery PS, Wang M, Xu C. Predictors of health-related quality of life in patients with symptomatic peripheral arterial disease. *J Vasc Surg* 2018;**68**:1126–34. <https://doi.org/10.1016/j.jvs.2017.12.074>
 167. Erickson KI, Hillman C, Stillman CM, Ballard RM, Bloodgood B, Conroy DE, et al. Physical activity, cognition, and brain outcomes: a review of the 2018 physical activity guidelines. *Med Sci Sports Exerc* 2019;**51**:1242–51. <https://doi.org/10.1249/MSS.0000000000001936>
 168. Kirk-Sanchez NJ, McGough EL. Physical exercise and cognitive performance in the elderly: current perspectives. *Clin Interv Aging* 2014;**9**:51–62. <https://doi.org/10.2147/CLIA.S39506>
 169. Gomez-Pinilla F, Hillman C. The influence of exercise on cognitive abilities. *Compr Physiol* 2013;**3**:403–28. <https://doi.org/10.1002/cphy.c110063>
 170. Gommans LN, Saarloos R, Scheltinga MR, Houterman S, de Bie RA, Fokkenrood HJ, et al. Editor's choice—the effect of supervision on walking distance in patients with intermittent claudication: a meta-analysis. *Eur J Vasc Endovasc Surg* 2014;**48**:169–84. <https://doi.org/10.1016/j.ejvs.2014.04.019>
 171. Abaraogu UO, Dall PM, Seenan CA. The effect of structured patient education on physical activity in patients with peripheral arterial disease and intermittent claudication: a systematic review. *Eur J Vasc Endovasc Surg* 2017;**54**:58–68. <https://doi.org/10.1016/j.ejvs.2017.04.003>
 172. Bearne LM, Volkmer B, Peacock J, Sekhon M, Fisher G, Galea Holmes MN, et al. Effect of a home-based, walking exercise behavior change intervention vs usual care on walking in adults with peripheral artery disease: the MOSAIC randomized clinical trial. *JAMA* 2022;**327**:1344–55. <https://doi.org/10.1001/jama.2022.3391>
 173. McDermott MM, Liu K, Guralnik JM, Criqui MH, Spring B, Tian L, et al. Home-based walking exercise intervention in peripheral artery disease: a randomized clinical trial. *JAMA* 2013;**310**:57–65. <https://doi.org/10.1001/jama.2013.7231>
 174. McDermott MM, Spring B, Berger JS, Treat-Jacobson D, Conte MS, Creager MA, et al. Effect of a home-based exercise intervention on peripheral artery disease: the HONOR randomized clinical trial. *JAMA* 2018;**319**:1665–76. <https://doi.org/10.1001/jama.2018.3275>
 175. Behrendt CA, Thomalla G, Rimmel DL, Petersen EL, Twerenbold R, Debus ES, et al. Prevalence of peripheral arterial disease, abdominal aortic aneurysm, and risk factors in the Hamburg City Health Study: a cross-sectional analysis. *Eur J Vasc Endovasc Surg* 2023;**65**:590–8. <https://doi.org/10.1016/j.ejvs.2023.01.002>
 176. Pabon M, Cheng S, Altin SE, Sethi SS, Nelson MD, Moreau KL, et al. Sex differences in peripheral artery disease. *Circ Res* 2022;**130**:496–511. <https://doi.org/10.1161/CIRCRESAHA.121.320702>
 177. Behrendt CA, Sigvant B, Kuchenbecker J, Grima MJ, Schermerhorn M, Thomson IA, et al. Editor's choice—international variations and sex disparities in the treatment of peripheral arterial occlusive disease: a report from VASCUNET and the International Consortium of Vascular Registries. *Eur J Vasc Endovasc Surg* 2020;**60**:873–80. <https://doi.org/10.1016/j.ejvs.2020.08.027>
 178. Detriche G, Guedon A, Mohamedi N, Sellami O, Cheng C, Galloway A, et al. Women specific characteristics and 1-year outcome among patients hospitalized for peripheral artery disease: a monocentric cohort analysis in a tertiary center. *Front Cardiovasc Med* 2022;**9**:824466. <https://doi.org/10.3389/fcvm.2022.824466>
 179. Heidemann F, Kuchenbecker J, Peters F, Kotov A, Marschall U, L'Hoest H, et al. A health insurance claims analysis on the effect of female sex on long-term outcomes after peripheral endovascular interventions for symptomatic peripheral arterial occlusive disease. *J Vasc Surg* 2021;**74**:780–7 e7. <https://doi.org/10.1016/j.jvs.2021.01.066>
 180. Hirsch AT, Allison MA, Gomes AS, Corriere MA, Duval S, Ershow AG, et al. A call to action: women and peripheral artery disease: a scientific statement from the American Heart Association. *Circulation* 2012;**125**:1449–72. <https://doi.org/10.1161/CIR.0b013e31824c39ba>
 181. Kotov A, Heidemann F, Kuchenbecker J, Peters F, Marschall U, Acar L, et al. Sex disparities in long term outcomes after open surgery for chronic limb threatening ischaemia: a propensity score matched analysis of health insurance claims. *Eur J Vasc Endovasc Surg* 2021;**61**:423–9. <https://doi.org/10.1016/j.ejvs.2020.11.006>
 182. Singh N, Liu K, Tian L, Criqui MH, Guralnik JM, Ferrucci L, et al. Leg strength predicts mortality in men but not in women with peripheral arterial disease. *J Vasc Surg* 2010;**52**:624–31. <https://doi.org/10.1016/j.jvs.2010.03.066>
 183. McDermott MM, Ferrucci L, Liu K, Guralnik JM, Tian L, Kibbe M, et al. Women with peripheral arterial disease experience faster functional decline than men with peripheral arterial disease. *J Am Coll Cardiol* 2011;**57**:707–14. <https://doi.org/10.1016/j.jacc.2010.09.042>
 184. Gommans LN, Scheltinga MR, van Sambeek MR, Maas AH, Bendermacher BL, Teijink JA. Gender differences following supervised exercise therapy in patients with intermittent claudication. *J Vasc Surg* 2015;**62**:681–8. <https://doi.org/10.1016/j.jvs.2015.03.076>
 185. Regensteiner JG, Bauer TA, Reusch JE, Brandenburg SL, Sippel JM, Vogelsong AM, et al. Abnormal oxygen uptake kinetic responses in women with type II diabetes mellitus. *J Appl Physiol* (1985) 1998;**85**:310–7. <https://doi.org/10.1152/jappl.1998.85.1.310>
 186. Lanzi S, Pousaz A, Calanca L, Mazzolai L. Sex-based differences in supervised exercise therapy outcomes for symptomatic peripheral artery disease. *Vasc Med* 2023;**28**:147–9. <https://doi.org/10.1177/1358863X221149454>
 187. Cetlin MD, Polonsky T, Ho K, Zhang D, Tian L, Zhao L, et al. Barriers to participation in supervised exercise therapy reported by people with peripheral artery disease. *J Vasc Surg* 2023;**77**:506–14. <https://doi.org/10.1016/j.jvs.2022.09.014>
 188. Gupta T, Manning P, Kolte D, Smolderen KG, Stone N, Henry JG, et al. Exercise therapy referral and participation in patients with peripheral artery disease: insights from the PORTRAIT registry. *Vasc Med* 2021;**26**:654–6. <https://doi.org/10.1177/1358863X211033649>
 189. Harwood A, Smith G, Broadbent E, Cayton T, Carradice D, Chetter I. Access to supervised exercise services for peripheral vascular disease patients. *Bull R Coll Surgeons Engl* 2017;**99**:207–11. <https://doi.org/10.1308/rcsbullet.2017.207>
 190. Harwood AE, Smith GE, Cayton T, Broadbent E, Chetter IC. A systematic review of the uptake and adherence rates to supervised exercise programs in patients with intermittent claudication. *Ann Vasc Surg* 2016;**34**:280–9. <https://doi.org/10.1016/j.avsg.2016.02.009>
 191. Li Y, Rother U, Rosenberg Y, Hinterseher I, Uhl C, Mylonas S, et al. A prospective survey study on the education and awareness about walking exercise amongst inpatients

- with symptomatic peripheral arterial disease in Germany. *Vasa* 2023;**52**:218–23. <https://doi.org/10.1024/0301-1526/a001057>
192. Rother U, Dorr G, Malyar N, Muller OJ, Steinbauer M, Ito W, et al. How German vascular surgeons and angiologists judge walking exercise for patients with PAD. *Vasa* 2023;**52**:224–9. <https://doi.org/10.1024/0301-1526/a001071>
193. Saxon JT, Safley DM, Mena-Hurtado C, Heyligers J, Fitridge R, Shishehbor M, et al. Adherence to guideline-recommended therapy-including supervised exercise therapy referral-across peripheral artery disease specialty clinics: insights from the International PORTRAIT Registry. *J Am Heart Assoc* 2020;**9**:e012541. <https://doi.org/10.1161/JAHA.119.012541>
194. Lanzi S, Belch J, Brodmann M, Madaric J, Bura-Riviere A, Visona A, et al. Supervised exercise training in patients with lower extremity peripheral artery disease. *Vasa* 2022;**51**:267–74. <https://doi.org/10.1024/0301-1526/a001024>
195. Parodi JC, Fernandez S, Moscovich F, Pulmaria C. Hydration may reverse most symptoms of lower extremity intermittent claudication or rest pain. *J Vasc Surg* 2020;**72**:1459–63. <https://doi.org/10.1016/j.jvs.2020.05.066>
196. Mannarino E, Pasqualini L, Innocente S, Scricciolo V, Rignanese A, Ciuffetti G. Physical training and antiplatelet treatment in stage II peripheral arterial occlusive disease: alone or combined? *Angiology* 1991;**42**:513–21. <https://doi.org/10.1177/000331979104200701>
197. Hobbs SD, Marshall T, Fegan C, Adam DJ, Bradbury AW. The effect of supervised exercise and cilostazol on coagulation and fibrinolysis in intermittent claudication: a randomized controlled trial. *J Vasc Surg* 2007;**45**:65–70; discussion 70 <https://doi.org/10.1016/j.jvs.2006.08.084>